

Anti-RPA2 Antibody (2Z862)

Product Details

Ig Type:	IgG
Reactivity:	Human,Mouse,Rat
Molecular Weight:	Theoretical: 29 kDa. Actual: 32 kDa.
Clone:	2Z862
Purification:	Protein A purified

Applications

Verified Activity:

1. 25 µg total protein per lane of various lysates (see on figure) probed with RPA2 monoclonal antibody, unconjugated (TMAB-12348) at 1:1000 dilution and 4°C overnight incubation. Followed by conjugated secondary antibody incubation at r. T. for 60 min.
2. Paraformaldehyde-fixed, paraffin embedded Mouse Liver; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with RPA2 Monoclonal Antibody, Unconjugated (TMAB-12348) at 1: 200 overnight at 4°C, followed by conjugation to the Goat Anti-Rabbit IgG H&L Secondary Antibody-HRP and DAB staining.
3. Paraformaldehyde-fixed, paraffin embedded Rat Liver; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with RPA2 Monoclonal Antibody, Unconjugated (TMAB-12348) at 1: 200 overnight at 4°C, followed by conjugation to the Goat Anti-Rabbit IgG H&L Secondary Antibody-HRP and DAB staining.
4. Paraformaldehyde-fixed, paraffin embedded Human Liver; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with RPA2 Monoclonal Antibody, Unconjugated (TMAB-12348) at 1: 200 overnight at 4°C, followed by conjugation to the Goat Anti-Rabbit IgG H&L Secondary Antibody-HRP and DAB staining.
5. Paraformaldehyde-fixed, paraffin embedded Mouse Cerebrum; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with RPA2 Monoclonal Antibody, Unconjugated (TMAB-12348) at 1: 200 overnight at 4°C, followed by conjugation to the Goat Anti-Rabbit IgG H&L Secondary Antibody-HRP and DAB staining.
6. Paraformaldehyde-fixed, paraffin embedded Rat Cerebrum; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with RPA2 Monoclonal Antibody, Unconjugated (TMAB-12348) at 1: 200 overnight at 4°C, followed by conjugation to the Goat Anti-Rabbit IgG H&L Secondary Antibody-HRP and DAB staining.
7. Paraformaldehyde-fixed, paraffin embedded Human Glioma; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with RPA2 Monoclonal Antibody, Unconjugated (TMAB-12348) at 1: 200 overnight at 4°C, followed by conjugation to the Goat Anti-Rabbit IgG H&L Secondary Antibody-HRP and DAB staining.
8. Paraformaldehyde-fixed, paraffin embedded Human Colon Cancer; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with RPA2 Monoclonal Antibody, Unconjugated (TMAB-12348) at 1: 200 overnight at 4°C, followed by conjugation to the Goat Anti-Rabbit IgG H&L Secondary Antibody-HRP and DAB staining.
9. Paraformaldehyde-fixed, paraffin embedded Human Prostate Cancer; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with RPA2 Monoclonal Antibody, Unconjugated (TMAB-12348) at 1: 200 overnight at 4°C, followed by conjugation to the Goat Anti-Rabbit IgG H&L Secondary Antibody-HRP and DAB staining.
10. Paraformaldehyde-fixed, paraffin embedded Human Breast Cancer; Antigen retrieval by

boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with RPA2 Monoclonal Antibody, Unconjugated (TMAB-12348) at 1: 200 overnight at 4°C, followed by conjugation to the Goat Anti-Rabbit IgG H&L Secondary Antibody-HRP and DAB staining.

11. Paraformaldehyde-fixed, paraffin embedded Human Colon; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with RPA2 Monoclonal Antibody, Unconjugated (TMAB-12348) at 1: 200 overnight at 4°C, followed by conjugation to the Goat Anti-Rabbit IgG H&L Secondary Antibody-HRP and DAB staining.

12. 4% Paraformaldehyde-fixed Hela (H) cell; Triton X-100 at r. T. for 20 min; Antibody incubation with (RPA2) monoclonal Antibody, unconjugated (TMAB-12348) 1:100, 90 min at 37°C; followed by conjugated Goat Anti-Rabbit IgG antibody (green) at 37°C for 90 min, DAPI (blue) was used to stain the cell nuclei. PBS instead of the primary antibody was used as the blank control.

Application: WB,IHC-P,IHC-Fr,ICC/IF,IF,FCM

Recommended WB: 1:500-2000; IHC-P: 1:100-500; IHC-Fr: 1:100-500; ICC/IF: 1:50-200; IF: 1:100-500; FCM: 1:50-100

Properties

Stability & Storage: Store at 2°C-8°C for 1 month. Store at -20°C or -80°C for 12 months. Avoid repeated freeze-thaw cycles.

Shipping: Shipping with blue ice.

Antigen Details

Gene ID: 6118

Research Background

Required for DNA recombination, repair and replication. The activity of RP-A is mediated by single-stranded DNA binding and protein interactions.

Functions as component of the alternative replication protein A complex (aRPA). aRPA binds single-stranded DNA and probably plays a role in DNA repair; it does not support chromosomal DNA replication and cell cycle progression through S-phase. In vitro, aRPA cannot promote efficient priming by DNA polymerase alpha but supports DNA polymerase delta synthesis in the presence of PCNA and replication factor C (RFC), the dual incision/excision reaction of nucleotide excision repair and RAD51-dependent strand exchange.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481