

Anti-SMC3 Antibody (9L985)

Product Details

Ig Type:	IgG
Reactivity:	Human,Mouse,Rat
Molecular Weight:	Theoretical: 142 kDa. Actual: 141 kDa.
Clone:	9L985
Purification:	Protein A purified

Applications

Verified Activity:

1. Paraformaldehyde-fixed, paraffin embedded Mouse Colon; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with SMC3 Monoclonal Antibody, Unconjugated (TMAB-12965) at 1:200 overnight at 4°C, followed by conjugation to the Goat Anti-Rabbit IgG H&L-HRP and DAB staining.
2. Paraformaldehyde-fixed, paraffin embedded Rat Colon; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with SMC3 Monoclonal Antibody, Unconjugated (TMAB-12965) at 1:200 overnight at 4°C, followed by conjugation to the Goat Anti-Rabbit IgG H&L-HRP and DAB staining.
3. Paraformaldehyde-fixed, paraffin embedded Human Colon; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with SMC3 Monoclonal Antibody, Unconjugated (TMAB-12965) at 1:200 overnight at 4°C, followed by conjugation to the Goat Anti-Rabbit IgG H&L-HRP and DAB staining.
4. Paraformaldehyde-fixed, paraffin embedded Mouse Stomach; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with SMC3 Monoclonal Antibody, Unconjugated (TMAB-12965) at 1:200 overnight at 4°C, followed by conjugation to the Goat Anti-Rabbit IgG H&L-HRP and DAB staining.
5. Paraformaldehyde-fixed, paraffin embedded Rat Stomach; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with SMC3 Monoclonal Antibody, Unconjugated (TMAB-12965) at 1:200 overnight at 4°C, followed by conjugation to the Goat Anti-Rabbit IgG H&L-HRP and DAB staining.
6. Paraformaldehyde-fixed, paraffin embedded Human Stomach; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with SMC3 Monoclonal Antibody, Unconjugated (TMAB-12965) at 1:200 overnight at 4°C, followed by conjugation to the Goat Anti-Rabbit IgG H&L-HRP and DAB staining.
7. Paraformaldehyde-fixed, paraffin embedded Mouse Cerebrum; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with SMC3 Monoclonal Antibody, Unconjugated (TMAB-12965) at 1:200 overnight at 4°C, followed by conjugation to the Goat Anti-Rabbit IgG H&L-HRP and DAB staining.
8. Paraformaldehyde-fixed, paraffin embedded Rat Cerebrum; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with SMC3 Monoclonal Antibody, Unconjugated (TMAB-12965) at 1:200 overnight at 4°C, followed by conjugation to the Goat Anti-Rabbit IgG H&L-HRP and DAB staining.
9. The HeLa (H) cells were fixed with 4% PFA (10 min at r. T.) and then permeabilized with 90% ice-cold methanol for 20 min at -20°C, the cells then were incubated in 5% BSA to block non-specific protein-protein interactions (30 min at r. T.), followed by secondary antibody incubation for 40 min at room temperature. Primary Antibody (green): Rabbit Anti-SMC3

A DRUG SCREENING EXPERT

antibody (TMAB-12965, 1: 100); Isotype Control (orange): Rabbit IgG. Blank control (black): PBS. Acquisition of 20,000 events was performed.

10. 4% Paraformaldehyde-fixed HeLa (H) cell; Triton X-100 at r. T. for 20 min; Antibody incubation with (SMC3) monoclonal Antibody, unconjugated (TMAB-12965) 1: 100, 90 min at 37°C; followed by conjugated Goat Anti-Rabbit IgG antibody (green, BF488) at 37°C for 90 min, DAPI (blue) was used to stain the cell nuclei. PBS instead of the primary antibody was used as the blank control.

Application: IHC-P,IHC-Fr,ICC/IF,IF,FCM

Recommended IHC-P: 1:100-500; IHC-Fr: 1:100-500; ICC/IF: 1:50-200; IF: 1:100-500; FCM: 1:50-100

Properties

Stability & Storage: Store at 2°C-8°C for 1 month. Store at -20°C or -80°C for 12 months. Avoid repeated freeze-thaw cycles.

Shipping: Shipping with blue ice.

Antigen Details

Immunogen: A synthesized peptide: human SMC3

Antigen Species: Human

Gene ID: 9126

Uniprot ID: Q9UQE7

Research Background

Central component of cohesin, a complex required for chromosome cohesion during the cell cycle. The cohesin complex may form a large proteinaceous ring within which sister chromatids can be trapped. At anaphase, the complex is cleaved and dissociates from chromatin, allowing sister chromatids to segregate. Cohesion is coupled to DNA replication and is involved in DNA repair. The cohesin complex also plays an important role in spindle pole assembly during mitosis and in chromosomes movement.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481
