

Anti-smooth muscle Myosin heavy chain 11 Antibody (2P75)

Product Details

Ig Type:	IgG
Reactivity:	Human,Mouse,Rat
Molecular Weight:	Theoretical: 227 kDa. Actual: 245 kDa.
Clone:	2P75
Purification:	Protein A purified

Applications

Verified Activity:

1. Paraformaldehyde-fixed, paraffin embedded Mouse Colon; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation with smooth muscle Myosin heavy chain 11 Monoclonal Antibody, Unconjugated (TMAB-12980) at 1: 200 overnight at 4°C, followed by conjugation to the Goat Anti-Rabbit IgG H&L Secondary Antibody-HRP and DAB staining.
2. Paraformaldehyde-fixed, paraffin embedded Rat Colon; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation with smooth muscle Myosin heavy chain 11 Monoclonal Antibody, Unconjugated (TMAB-12980) at 1: 200 overnight at 4°C, followed by conjugation to the Goat Anti-Rabbit IgG H&L Secondary Antibody-HRP and DAB staining.
3. Paraformaldehyde-fixed, paraffin embedded Human Colon; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation with smooth muscle Myosin heavy chain 11 Monoclonal Antibody, Unconjugated (TMAB-12980) at 1: 200 overnight at 4°C, followed by conjugation to the Goat Anti-Rabbit IgG H&L Secondary Antibody-HRP and DAB staining.
4. Paraformaldehyde-fixed, paraffin embedded Mouse Prostate; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation with smooth muscle Myosin heavy chain 11 Monoclonal Antibody, Unconjugated (TMAB-12980) at 1: 200 overnight at 4°C, followed by conjugation to the Goat Anti-Rabbit IgG H&L Secondary Antibody-HRP and DAB staining.
5. Paraformaldehyde-fixed, paraffin embedded Human Prostate; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation with smooth muscle Myosin heavy chain 11 Monoclonal Antibody, Unconjugated (TMAB-12980) at 1: 200 overnight at 4°C, followed by conjugation to the Goat Anti-Rabbit IgG H&L Secondary Antibody-HRP and DAB staining.
6. Paraformaldehyde-fixed, paraffin embedded Mouse Stomach; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation with smooth muscle Myosin heavy chain 11 Monoclonal Antibody, Unconjugated (TMAB-12980) at 1: 200 overnight at 4°C, followed by conjugation to the Goat Anti-Rabbit IgG H&L Secondary Antibody-HRP and DAB staining.
7. Paraformaldehyde-fixed, paraffin embedded Rat Stomach; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation with smooth muscle Myosin heavy chain 11 Monoclonal Antibody, Unconjugated (TMAB-12980) at 1: 200 overnight at 4°C, followed by conjugation to the Goat Anti-Rabbit IgG H&L Secondary Antibody-HRP and DAB staining.
8. Paraformaldehyde-fixed, paraffin embedded Human Stomach; Antigen retrieval by boiling

in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation with smooth muscle Myosin heavy chain 11 Monoclonal Antibody, Unconjugated (TMAB-12980) at 1: 200 overnight at 4°C, followed by conjugation to the Goat Anti-Rabbit IgG H&L Secondary Antibody-HRP and DAB staining.

9. Paraformaldehyde-fixed, paraffin embedded Mouse Uterus; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation with smooth muscle Myosin heavy chain 11 Monoclonal Antibody, Unconjugated (TMAB-12980) at 1: 200 overnight at 4°C, followed by conjugation to the Goat Anti-Rabbit IgG H&L Secondary Antibody-HRP and DAB staining.

10. Paraformaldehyde-fixed, paraffin embedded Rat Uterus; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation with smooth muscle Myosin heavy chain 11 Monoclonal Antibody, Unconjugated (TMAB-12980) at 1: 200 overnight at 4°C, followed by conjugation to the Goat Anti-Rabbit IgG H&L Secondary Antibody-HRP and DAB staining.

11. Paraformaldehyde-fixed, paraffin embedded Human Uterus; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation with smooth muscle Myosin heavy chain 11 Monoclonal Antibody, Unconjugated (TMAB-12980) at 1: 200 overnight at 4°C, followed by conjugation to the Goat Anti-Rabbit IgG H&L Secondary Antibody-HRP and DAB staining.

12. Paraformaldehyde-fixed, paraffin embedded Human Lung Cancer; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation with smooth muscle Myosin heavy chain 11 Monoclonal Antibody, Unconjugated (TMAB-12980) at 1: 200 overnight at 4°C, followed by conjugation to the Goat Anti-Rabbit IgG H&L Secondary Antibody-HRP and DAB staining.

13. Paraformaldehyde-fixed, paraffin embedded Human Placenta; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation with smooth muscle Myosin heavy chain 11 Monoclonal Antibody, Unconjugated (TMAB-12980) at 1: 200 overnight at 4°C, followed by conjugation to the Goat Anti-Rabbit IgG H&L Secondary Antibody-HRP and DAB staining.

14. Paraformaldehyde-fixed, paraffin embedded Human Gastric Cancer; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation with smooth muscle Myosin heavy chain 11 Monoclonal Antibody, Unconjugated (TMAB-12980) at 1: 200 overnight at 4°C, followed by conjugation to the Goat Anti-Rabbit IgG H&L Secondary Antibody-HRP and DAB staining.

15. 25 µg total protein per lane of various lysates (see on figure) probed with smooth muscle Myosin heavy chain 11 monoclonal antibody, unconjugated (TMAB-12980) at 1:1000 dilution and 4°C overnight incubation. Followed by conjugated secondary antibody incubation at r. T. for 60 min.

Application: WB,IHC-P,IF,IHC-Fr

Recommended WB: 1:500-2000; IHC-P: 1:100-500; IF: 1:100-500; IHC-Fr: 1:100-500

Properties

Stability & Storage: Store at 2°C-8°C for 1 month. Store at -20°C or -80°C for 12 months. Avoid repeated freeze-thaw cycles.

Shipping: Shipping with blue ice.

Antigen Details

Immunogen:	A synthesized peptide: human Myosin heavy chain 11
Antigen Species:	Human
Gene ID:	4629
Uniprot ID:	P35749

Research Background

The protein encoded by this gene is a smooth muscle myosin belonging to the myosin heavy chain family. The gene product is a subunit of a hexameric protein that consists of two heavy chain subunits and two pairs of non-identical light chain subunits. It functions as a major contractile protein, converting chemical energy into mechanical energy through the hydrolysis of ATP. The gene encoding a human ortholog of rat NUDE1 is transcribed from the reverse strand of this gene, and its 3' end overlaps with that of the latter. The pericentric inversion of chromosome 16[inv(16)(p13q22)] produces a chimeric transcript that encodes a protein consisting of the first 165 residues from the N terminus of core-binding factor beta in a fusion with the C-terminal portion of the smooth muscle myosin heavy chain. This chromosomal rearrangement is associated with acute myeloid leukemia of the M4Eo subtype. Alternative splicing generates isoforms that are differentially expressed, with ratios changing during muscle cell maturation. Alternatively spliced transcript variants encoding different isoforms have been identified.

MYH11 is a smooth muscle myosin belonging to the myosin heavy chain family. It is a subunit of a hexameric protein that consists of two heavy chain subunits and two pairs of non-identical light chain subunits. It functions as a major contractile protein, converting chemical energy into mechanical energy through the hydrolysis of ATP.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481