

Anti-TARDBP Antibody (4W355)

Product Details

Ig Type:	IgG
Reactivity:	Human,Mouse,Rat (predicted:Zebrafish)
Molecular Weight:	Theoretical: 45 kDa. Actual: 45 kDa.
Clone:	4W355
Purification:	Protein A purified

Applications

Verified Activity:

1. Paraformaldehyde-fixed, paraffin embedded Mouse Spinal Cord; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation TARDBP Monoclonal Antibody, Unconjugated (TMAB-13381) at 1:200 overnight at 4°C, followed by conjugation to the Goat Anti-Rabbit IgG H&L-HRP and DAB staining.
2. Paraformaldehyde-fixed, paraffin embedded Rat Spinal Cord; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation TARDBP Monoclonal Antibody, Unconjugated (TMAB-13381) at 1:200 overnight at 4°C, followed by conjugation to the Goat Anti-Rabbit IgG H&L-HRP and DAB staining.
3. Paraformaldehyde-fixed, paraffin embedded Mouse Colon; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation TARDBP Monoclonal Antibody, Unconjugated (TMAB-13381) at 1:200 overnight at 4°C, followed by conjugation to the Goat Anti-Rabbit IgG H&L-HRP and DAB staining.
4. Paraformaldehyde-fixed, paraffin embedded Rat Colon; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation TARDBP Monoclonal Antibody, Unconjugated (TMAB-13381) at 1:200 overnight at 4°C, followed by conjugation to the Goat Anti-Rabbit IgG H&L-HRP and DAB staining.
5. Paraformaldehyde-fixed, paraffin embedded Human Colon; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation TARDBP Monoclonal Antibody, Unconjugated (TMAB-13381) at 1:200 overnight at 4°C, followed by conjugation to the Goat Anti-Rabbit IgG H&L-HRP and DAB staining.
6. Paraformaldehyde-fixed, paraffin embedded Mouse Cerebrum; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation TARDBP Monoclonal Antibody, Unconjugated (TMAB-13381) at 1:200 overnight at 4°C, followed by conjugation to the Goat Anti-Rabbit IgG H&L-HRP and DAB staining.
7. Paraformaldehyde-fixed, paraffin embedded Rat Cerebrum; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation TARDBP Monoclonal Antibody, Unconjugated (TMAB-13381) at 1:200 overnight at 4°C, followed by conjugation to the Goat Anti-Rabbit IgG H&L-HRP and DAB staining.
8. Paraformaldehyde-fixed, paraffin embedded Human Cerebrum; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation TARDBP Monoclonal Antibody, Unconjugated (TMAB-13381) at 1:200 overnight at 4°C, followed by conjugation to the Goat Anti-Rabbit IgG H&L-HRP and DAB staining.
9. Paraformaldehyde-fixed, paraffin embedded Mouse Cerebellum; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation TARDBP Monoclonal Antibody, Unconjugated (TMAB-13381) at 1:200 overnight at 4°C, followed by conjugation to the Goat Anti-Rabbit IgG H&L-HRP and DAB staining.

10. Paraformaldehyde-fixed, paraffin embedded Rat Cerebellum; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation TARDBP Monoclonal Antibody, Unconjugated (TMAB-13381) at 1:200 overnight at 4°C, followed by conjugation to the Goat Anti-Rabbit IgG H&L-HRP and DAB staining.

11. Paraformaldehyde-fixed, paraffin embedded Human Glioma; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation TARDBP Monoclonal Antibody, Unconjugated (TMAB-13381) at 1:200 overnight at 4°C, followed by conjugation to the Goat Anti-Rabbit IgG H&L-HRP and DAB staining.

12. Paraformaldehyde-fixed, paraffin embedded Human Colon Cancer; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation TARDBP Monoclonal Antibody, Unconjugated (TMAB-13381) at 1:200 overnight at 4°C, followed by conjugation to the Goat Anti-Rabbit IgG H&L-HRP and DAB staining.

13. Paraformaldehyde-fixed, paraffin embedded Human Prostate Cancer; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation TARDBP Monoclonal Antibody, Unconjugated (TMAB-13381) at 1:200 overnight at 4°C, followed by conjugation to the Goat Anti-Rabbit IgG H&L-HRP and DAB staining.

14. Paraformaldehyde-fixed, paraffin embedded Human Endometrial Cancer; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation TARDBP Monoclonal Antibody, Unconjugated (TMAB-13381) at 1:200 overnight at 4°C, followed by conjugation to the Goat Anti-Rabbit IgG H&L-HRP and DAB staining.

15. 25 ug total protein per lane of various lysates (see on figure) probed with TARDBP monoclonal antibody, unconjugated (TMAB-13381) at 1:2000 dilution and 4°C overnight incubation. Followed by conjugated secondary antibody incubation at r. T. for 60 min.

16. Paraformaldehyde-fixed, paraffin embedded Rat Cerebrum; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with TARDBP Monoclonal Antibody, Unconjugated (TMAB-13381) at 1:200 overnight at 4°C. Followed by conjugated Goat Anti-Rabbit IgG antibody (Red, Goat Anti-Rabbit IgG H&L-BF594), DAPI (blue) was used to stain the cell nuclei.

17. Paraformaldehyde-fixed, paraffin embedded Human Cerebrum; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with TARDBP Monoclonal Antibody, Unconjugated (TMAB-13381) at 1:200 overnight at 4°C. Followed by conjugated Goat Anti-Rabbit IgG antibody (Red, Goat Anti-Rabbit IgG H&L-BF594), DAPI (blue) was used to stain the cell nuclei.

18. Paraformaldehyde-fixed, paraffin embedded Mouse Cerebrum; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with TARDBP Monoclonal Antibody, Unconjugated (TMAB-13381) at 1:200 overnight at 4°C. Followed by conjugated Goat Anti-Rabbit IgG antibody (Red, Goat Anti-Rabbit IgG H&L-BF594), DAPI (blue) was used to stain the cell nuclei.

Application: WB,IHC-P,IHC-Fr,ICC/IF,IF,FCM

Recommended WB: 1:1000-5000; IHC-P: 1:100-500; IHC-Fr: 1:100-500; ICC/IF: 1:100-500; IF: 1:100-500; FCM: 1:50-100

Properties

Stability & Storage: Store at 2°C-8°C for 1 month. Store at -20°C or -80°C for 12 months. Avoid repeated freeze-thaw cycles.

Shipping: Shipping with blue ice.

Antigen Details

Immunogen: A synthesized peptide: human TDP43
Antigen Species: Human
Gene ID: 23435
Uniprot ID: Q13148

Research Background

HIV-1, the causative agent of acquired immunodeficiency syndrome (AIDS), contains an RNA genome that produces a chromosomally integrated DNA during the replicative cycle. Activation of HIV-1 gene expression by the transactivator Tat is dependent on an RNA regulatory element (TAR) located downstream of the transcription initiation site. The protein encoded by this gene is a transcriptional repressor that binds to chromosomally integrated TAR DNA and represses HIV-1 transcription. In addition, this protein regulates alternate splicing of the CFTR gene. A similar pseudogene is present on chromosome 20. [provided by RefSeq, Jul 2008]

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481