

Anti-VP16 tag Polyclonal Antibody

Product Details

Ig Type:	IgG
Reactivity:	Species independent
Molecular Weight:	Actual: 51 kDa.
Purification:	Protein A purified

Applications

Sample:	Lane 1: VP16-Tagged Fusion Protein Overexpression E. Coli Lysate at 2 µg Lane 2: VP16-Tagged Fusion Protein Overexpression E. Coli Lysate at 4 µg
Verified Activity:	Primary: Anti-VP16 tag (TMAB-14126) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 51 kD Observed band size: 51 kD
Application:	ELISA,FCM,ICC/IF,IF,IHC-Fr,IHC-P,WB
Recommended	ELISA=1:1000-5000; FCM=1 µg/Test; ICC/IF=1:100-500; IF=1:200-1000; IHC-Fr=1:200-1000; IHC-P=1:200-1000; WB=1:1000-5000

Properties

Stability & Storage:	Store at 2°C-8°C for 1 month. Store at -20°C or -80°C for 12 months. Avoid repeated freeze-thaw cycles.
Shipping:	Shipping with blue ice.

Antigen Details

Immunogen:	KLH conjugated synthetic peptide: HSV-1 VP16
------------	--

Research Background

HSV evades the immune system through interference with MHC class I antigen presentation on the cell surface, by blocking TAP or the transporter associated with antigen processing induced by the secretion of ICP-47 by HSV. In the host cell, TAP transports digested viral antigen epitope peptides from the cytosol to the endoplasmic reticulum, allowing these epitopes to be combined with MHC class I molecules and presented on the surface of the cell. Viral epitope presentation with MHC class I is a requirement for activation of cytotoxic T-lymphocytes (CTLs), the major effectors of the cell-mediated immune response against virally-infected cells. ICP-47 prevents initiation of a CTL-response against HSV, allowing the virus to survive for a protracted period in the host.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only. Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481