


Chebulagic acid

Chemical Properties

CAS No. :	23094-71-5
Formula:	C ₄₁ H ₃₀ O ₂₇
Molecular Weight:	954.66
Storage:	Keep away from direct sunlight Powder: -20°C for 3 years In solvent: -80°C for 1 year <i>Actual storage temperature shall be subject to the COA.</i>



Biological Description

Description	Chebulagic acid, isolated from the Terminalia chebula Retz, is a COX-LOX dual inhibitor.
Targets(IC50)	COX, Influenza Virus, Lipoyxygenase, SARS-CoV
In vitro	Chebulagic acid can enhance autophagy and exerts anti-inflammatory and anti-infective effects [1,2]. Chebulagic acid also shows a protective effect against 1-methyl-4-phenylpyridinium (MPP+) - induce cytotoxicity which mimics the pathological symptom of Parkinson's disease. Chebulagic acid inhibits the LPS-induced upregulation of TNF- α and IL-1 β in a dose- and time-dependent manner [3].

Solubility Information

Solubility	DMSO: 150 mg/mL (157.12 mM), Sonication is recommended. (< 1 mg/ml refers to the product slightly soluble or insoluble)
In vivo Formulation	10% DMSO+90% Saline: 10 mg/mL (10.47 mM), Solution. 10% DMSO+40% PEG300+5% Tween-80+45% Saline: 1 mg/mL (1.05 mM), Sonication is recommended. <i>Please add the solvents sequentially, clarifying the solution as much as possible before adding the next one. Dissolve by heating and/or sonication if necessary. Working solution is recommended to be prepared and used immediately. The formulation provided above is for reference purposes only. In vivo formulations may vary and should be modified based on specific experimental conditions.</i>

Preparing Stock Solutions

	1mg	5mg	10mg
1 mM	1.0475 mL	5.2375 mL	10.4749 mL
5 mM	0.2095 mL	1.0475 mL	2.095 mL
10 mM	0.1047 mL	0.5237 mL	1.0475 mL
50 mM	0.0209 mL	0.1047 mL	0.2095 mL

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. Please use it as soon as possible.

Note: The dilution table applies only to solid products. For liquid products, please calculate the stock solution based on the stated concentration and/or density.

Reference

Kim HJ et al. Neuroprotective Effect of Chebulagic Acid via Autophagy Induction in SH-SY5Y Cells. *Biomol Ther* (Seoul). 2014 Jul;22(4):275-81.

Liu Y et al. Chebulagic acid inhibits the LPS-induced expression of TNF- α and IL-1 β in endothelial cells by suppressing MAPK activation. *Exp Ther Med*. 2015 Jul;10(1):263-268.

Athira AP et al. Inhibition of Angiogenesis In Vitro by Chebulagic Acid: A COX-LOX Dual Inhibitor. *Int J Vasc Med*. 2013;2013:843897.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481