

## Integrin alpha 4 beta 7 Protein, Cynomolgus, Recombinant (His)

### General Information

Synonyms:	alpha 4;β7;LPAM;ITGA4&ITGB7;beta7;Integrin alpha 4 β 7;Integrin alpha 4 beta 7
Protein Construction:	Tyr34-Thr977 (ITGA4) acidic tail and Glu20-His723 (ITGB7) basic tail
Species:	Cynomolgus
Expression Host:	HEK293 Cells
Accession:	XP_005573683.3(ITGA4)&XP_005571028.3(ITGB7)
Molecular Weight:	110.87 kDa (ITGA4) and 81.09 kDa (ITGB7) (Predicted); 73-80 kDa, 110-120 kDa and 130-150 kDa (Reducing condition, due to glycosylation)

### QC Testing

Biological Activity:	Immobilized Cynomolgus ITGA4&ITGB7, His Tag at 2 µg/ml (100 µl/well) on the plate. Dose response curve for Anti-ITGA4 Antibody, hFc Tag with the EC50 of 16.7 ng/ml determined by ELISA (QC Test).
Purity:	> 95% as determined by Bis-Tris PAGE
Endotoxin:	< 1 EU/µg of the protein as determined by the LAL method.
Formulation:	Lyophilized from 0.22 µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.

### Preparation and Storage

#### Reconstitution:

Reconstitute the lyophilized protein in distilled water. The product concentration should not be less than 100 µg/ml. Before opening, centrifuge the tube to collect powder at the bottom. After adding the reconstitution buffer, avoid vortexing or pipetting for mixing.

#### Stability & Storage:

It is recommended to store recombinant proteins at -20°C to -80°C for future use. Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Actual storage temperature shall be subject to the COA.

#### Shipping:

In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

### Protein Background

Integrins alpha-4/beta-1 (VLA-4) and alpha-4/beta-7 are receptors for fibronectin. They recognize one or more domains within the alternatively spliced CS-1 and CS-5 regions of fibronectin. They are also receptors for VCAM1. Integrin alpha-4/beta-1 recognizes the sequence Q-I-D-S in VCAM1. Integrin alpha-4/beta-7 is also a receptor for MADCAM1. It recognizes the sequence L-D-T in MADCAM1.

**Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins**

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E\_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481