

A DRUG SCREENING EXPERT

In vivo Formulation	<i>vary and should be modified based on specific experimental conditions.</i>
---------------------	---

Preparing Stock Solutions

	1mg	5mg	10mg
1 mM	1.9606 mL	9.8032 mL	19.6063 mL
5 mM	0.3921 mL	1.9606 mL	3.9213 mL
10 mM	0.1961 mL	0.9803 mL	1.9606 mL
50 mM	0.0392 mL	0.1961 mL	0.3921 mL

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. Please use it as soon as possible.

Note: The dilution table applies only to solid products. For liquid products, please calculate the stock solution based on the stated concentration and/or density.

Reference

Obach RS, et al. Drug Metab Dispos, 2004, 32(1), 89-97.

Liu Z, Wang Y, Qin W, et al. Raloxifene alleviates amyloid- β -induced cytotoxicity in HT22 neuronal cells via inhibiting oligomeric and fibrillar species formation. Journal of Biochemical and Molecular Toxicology. 2019

Yang NN, et al. Endocrinology, 1996, 137(5), 2075-2084.

Black LJ, et al. J Clin Invest, 1994, 93(1), 63-69.

Evans G, et al. Endocrinology, 1994, 134(5), 2283-2288.

Sato M, et al. FASEB J, 1996, 10(8), 905-912.

Liu Z, Wang Y, Qin W, et al. Raloxifene alleviates amyloid- β -induced cytotoxicity in HT22 neuronal cells via inhibiting oligomeric and fibrillar species formation[J]. Journal of biochemical and molecular toxicology. 2019.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel: 781-999-4286 E_mail: info@targetmol.com Address: 34 Washington Street, Wellesley Hills, MA 02481