

Suc-Leu-Tyr-AMC

Chemical Properties

CAS No. : 94367-20-1

Formula: C₂₉H₃₃N₃O₈


Molecular Weight: 551.59

Storage:

Keep away from moisture, Store at low temperature,
Keep away from direct sunlight

Powder: -20°C for 3 years | In solvent: -80°C for 1 year

Actual storage temperature shall be subject to the COA.



Biological Description

Description	Suc-Leu-Tyr-AMC is a fluorescent substrate for calpain I and II, papain (another cysteine protease), and is used to measure the chymotrypsin-like peptidase activity of the 20S proteasome (excitation max: 360 nm; emission max: 460 nm). Suc-Leu-Tyr-AMC can also be cleaved by the Ti protease from E. coli.
Targets(IC50)	Others

Solubility Information

Solubility	DMF: 30 mg/mL (54.39 mM), Sonication is recommended. DMSO:PBS(pH7.2) (1:1): 0.5 mg/mL (0.91 mM), Sonication is recommended. Ethanol: 20 mg/mL (36.26 mM), Sonication is recommended. DMSO: 30 mg/mL (54.39 mM), Sonication is recommended. (< 1 mg/ml refers to the product slightly soluble or insoluble)
------------	---

Preparing Stock Solutions

	1mg	5mg	10mg
1 mM	1.8129 mL	9.0647 mL	18.1294 mL
5 mM	0.3626 mL	1.8129 mL	3.6259 mL
10 mM	0.1813 mL	0.9065 mL	1.8129 mL
50 mM	0.0363 mL	0.1813 mL	0.3626 mL

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. Please use it as soon as possible.

Note: The dilution table applies only to solid products. For liquid products, please calculate the stock solution based on the stated concentration and/or density.

Reference

Woo KM, et al. Protease Ti from Escherichia coli requires ATP hydrolysis for protein breakdown but not for hydrolysis of small peptides. J Biol Chem. 1989;264(4):2088-2091.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481