


## Buparlisib

## Chemical Properties

CAS No. :	944396-07-0
Formula:	C <sub>18</sub> H <sub>21</sub> F <sub>3</sub> N <sub>6</sub> O <sub>2</sub>
Molecular Weight:	410.39
Storage:	Powder: -20°C for 3 years   In solvent: -80°C for 1 year Actual storage temperature shall be subject to the COA.



## Biological Description

Description	Buparlisib (BKM120) is an orally bioavailable specific oral inhibitor of the pan-class I PI3K (IC50s: 52/166/116/nM for p110 $\alpha$ , p110 $\beta$ , and p110 $\delta$ ).
Targets(IC50)	Apoptosis,PI3K
In vitro	Buparlisib exhibited 50-300 nM activity for class I PI3K's, including the most common p110R mutants. Additionally, 15 exhibited lower potency against class III and class IV PI3K's, where 2, 5, >5, and >25 $\mu$ M biochemical activity was observed for inhibition of VPS34, mTOR, DNAPK, and PI4K, respectively. Across all cell lines, pathway modulation and antiproliferative activity was consistent with cellular PI3K inhibition [1]. Buparlisib demonstrated anti-proliferative activity in 11 human gastric cancer cell lines by decreasing mTOR downstream signaling. But Buparlisib treatment increased p-AKT by subsequent abrogation of feedback inhibition by stabilizing insulin receptor substrate-1. In KRAS mutant gastric cancer cells, either p-ERK or p-STAT3 was also increased upon treatment of Buparlisib [2]. BKM120 induces cell growth inhibition and apoptosis in both multiple myeloma (MM) cell lines and freshly isolated primary MM cells. In addition, BKM120 shows synergistic cytotoxicity with dexamethasone in dexamethasone-sensitive MM cells. Low doses of BKM120 and dexamethasone, each of which alone has limited cytotoxicity, induce significant cell apoptosis in MM.1S and ARP-1. BKM120 exposure causes cell cycle arrest by upregulating p27 (Kip1) and downregulating cyclin D1 and induces caspase-dependent apoptosis by downregulating antiapoptotic XIAP and upregulating expression of a cytotoxic small isoform of Bim, BimS [3].
In vivo	In A2780 xenograft tumors, oral administration of Buparlisib at 3, 10, 30, 60, and 100 mg/kg resulted in dose-dependent modulation of pAKTSer473. Partial inhibition of pAKTSer473 was observed at 3 and 10 mg/kg, with near complete inhibition at 30, 60, or 100 mg/kg. Inhibition of pAkt corresponded with plasma and tumor drug exposure and was time-dependent, with over 90% target modulation achieved at 60 and 100 mg/kg doses at the 10-hour mark when plasma and tumor exposure was approximately 2 $\mu$ M [1]. Buparlisib treatment (5 $\mu$ M/kg/day) significantly reduced tumor burdens and circulating human kappa chain levels, also extending the survival of tumor-bearing mice [3].
Kinase Assay	Compounds to be tested were dissolved in DMSO and directly distributed into a black 384-well plate at 1.25 $\mu$ L per well. To start the reaction, 25 $\mu$ L of 10 nM PI3 kinase and 5 $\mu$ g/mL 1-alpha-phosphatidylinositol (PI) in assay buffer (10 mM Tris pH 7.5, 5 mM

## A DRUG SCREENING EXPERT

Kinase Assay	MgCl <sub>2</sub> , 20 mM NaCl, 1 mM DTT and 0.05% CHAPS) were added into each well followed by 25 µL of 2 µM ATP in assay buffer. The reaction was performed until approx 50% of the ATP was depleted, and then stopped by the addition of 25 µL of KinaseGlo solution. The stopped reaction was incubated for 5 minutes and the remaining ATP was then detected via luminescence [1].
Cell Research	A2780 cells were cultured in DMEM supplemented with 10% FBS. L-glutamine, sodium pyruvate, and antibiotics. Cells were plated in the same medium at a density of 1000 cells per well, 100 ul per well into black-walled-clear-bottom plates and incubated for 3-5 hours. Test compounds supplied in DMSO (20 mM) were diluted further into DMSO (7.5 ul of 20 mM test compound in 22.5 ul DMSO. Mix well, transfer 10 ul to 20 ul DMSO, repeat until 9 concentrations have been made). The diluted test compound solution (2uL), was then added to the cell medium (500 ul) cell medium. Equal volumes of this solution (100 uL) were added to the cells in 96 well plates and incubated at 37 oC for 3 days and developed using Cell Titer Glo. Inhibition of cell proliferation was determined by luminescence read using Trilux [1].
Animal Research	Six- to eight-week-old female severe combined immunodeficiency (SCID) mice were housed and monitored in the MD Anderson Cancer Center animal research facility. SCID mice were subcutaneously inoculated in the right flank with 1 million ARP-1 or MM.1S cells suspended in 50 µl phosphate-buffered saline (PBS). After palpable tumor developed (tumor diameter ≥5 mm), mice were treated with intraperitoneal injection of DMSO/PBS or BKM120 (5 µM per kg per day) for 15 days. Tumor sizes were measured every 5 days, and blood samples were collected at the same period. Tumor burdens were evaluated by measuring tumor size and detecting the circulating human kappa chain or lambda chain [3].

### Solubility Information

Solubility	DMSO: 125 mg/mL (304.59 mM),Sonication is recommended. H <sub>2</sub> O: Insoluble, Ethanol: 2 mg/mL (4.87 mM),Sonication is recommended. (< 1 mg/ml refers to the product slightly soluble or insoluble)
In vivo Formulation	10% DMSO+40% PEG300+5% Tween 80+45% Saline: 10 mg/mL (24.37 mM),Suspension. 10% DMSO+90% Saline: < 10 mg/mL (24.37 mM),Lower concentrations may be soluble, but exact solubility limit is unknown. <i>Please add the solvents sequentially, clarifying the solution as much as possible before adding the next one. Dissolve by heating and/or sonication if necessary. Working solution is recommended to be prepared and used immediately. The formulation provided above is for reference purposes only. In vivo formulations may vary and should be modified based on specific experimental conditions.</i>

## Preparing Stock Solutions

	1mg	5mg	10mg
1 mM	2.4367 mL	12.1835 mL	24.3671 mL
5 mM	0.4873 mL	2.4367 mL	4.8734 mL
10 mM	0.2437 mL	1.2184 mL	2.4367 mL
50 mM	0.0487 mL	0.2437 mL	0.4873 mL

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. Please use it as soon as possible.

Note: The dilution table applies only to solid products. For liquid products, please calculate the stock solution based on the stated concentration and/or density.

## Reference

Burger MT, et al. Identification of NVP-BKM120 as a Potent, Selective, Orally Bioavailable Class I PI3 Kinase Inhibitor for Treating Cancer. ACS Med Chem Lett. 2011 Aug 26;2(10):774-9.

Wencao Zhao, Le Cao, Hanru Ying, Wenjuan Zhang, Dantong Li, Xiaolong Zhu, Wenzhi Xue, Shuang Wu, Mengye Cao, Cong Fu, Haonan Qi, Yimei Hao, Yun-Chi Tang, Jun Qin, Tao P. Zhong, Xiaoxi Lin, Luyang Yu, Xuri Li, Lin Li, Dianqing Wu & Weijun Pan Endothelial CDS2 deficiency causes VEGFA-mediated vascular regression and tumor inhibition. Cell Research. 2019

Hu P, Li H, Yu X, et al. GL-V9 exerts anti-T cell malignancies effects via promoting lysosome-dependent AKT1 degradation and activating AKT1/FOXO3A/BIM axis. Free Radical Biology and Medicine. 2019

Park E, et al. NVP-BKM120, a novel PI3K inhibitor, shows synergism with a STAT3 inhibitor in human gastric cancer cells harboring KRAS mutations. Int J Oncol. 2012 Apr;40(4):1259-66.

Zheng Y, et al. Novel phosphatidylinositol 3-kinase inhibitor NVP-BKM120 induces apoptosis in myeloma cells and shows synergistic anti-myeloma activity with dexamethasone. J Mol Med (Berl). 2012 Jun;90(6):695-706.

Hu P, Li H, Yu X, et al. GL-V9 exerts anti-T cell malignancies effects via promoting lysosome-dependent AKT1 degradation and activating AKT1/FOXO3A/BIM axis[J]. Free Radical Biology and Medicine. 2019.

Wencao Zhao, Le Cao, Hanru Ying, Wenjuan Zhang, Dantong Li, Xiaolong Zhu, Wenzhi Xue, Shuang Wu, Mengye Cao, Cong Fu, Haonan Qi, Yimei Hao, Yun-Chi Tang, Jun Qin, Tao P. Zhong, Xiaoxi Lin, Luyang Yu, Xuri Li, Lin Li, Dianqing Wu & Weijun Pan . Endothelial CDS2 deficiency causes VEGFA-mediated vascular regression and tumor inhibition. Cell Research. 2019

**Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins**

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E\_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481