


S-Dihydrodaidzein

Chemical Properties

CAS No. :	879559-75-8
Formula:	C ₁₅ H ₁₂ O ₄
Molecular Weight:	256.25
Storage:	Powder: -20°C for 3 years In solvent: -80°C for 1 year Actual storage temperature shall be subject to the COA.



Biological Description

Description	S-Dihydrodaidzein ((3S)-7-hydroxy-3-(4-hydroxyphenyl)-2,3-dihydrochromen-4-one) is an enantiomer of dihydrodaidzein which is involved in equol biosynthesis in a lactic acid bacterium, Lactococcus sp. strain.
Targets(IC50)	Others

Solubility Information

Solubility	DMSO: 250 mg/mL (975.61 mM), Sonication is recommended. (< 1 mg/ml refers to the product slightly soluble or insoluble)
------------	---

Preparing Stock Solutions

	1mg	5mg	10mg
1 mM	3.9024 mL	19.5122 mL	39.0244 mL
5 mM	0.7805 mL	3.9024 mL	7.8049 mL
10 mM	0.3902 mL	1.9512 mL	3.9024 mL
50 mM	0.078 mL	0.3902 mL	0.7805 mL

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. Please use it as soon as possible.

Note: The dilution table applies only to solid products. For liquid products, please calculate the stock solution based on the stated concentration and/or density.

Reference

- Shimada Y, et al. Identification of a novel dihydrodaidzein racemase essential for biosynthesis of equol from daidzein in Lactococcus sp. strain 20-92. Appl Environ Microbiol. 2012 Jul;78(14):4902-7.
- Franke AA, et al. Liquid chromatographic-photodiode array mass spectrometric analysis of dietary phytoestrogens from human urine and blood. J Chromatogr B Analyt Technol Biomed Life Sci. 2002 Sep 25;777(1-2):45-59.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481