


mTOR inhibitor-8

Chemical Properties

CAS No. :	2489196-70-3
Formula:	C ₂₄ H ₁₉ ClN ₄ O ₅
Molecular Weight:	446.95
Storage:	Powder: -20°C for 3 years In solvent: -80°C for 1 year Actual storage temperature shall be subject to the COA.



Biological Description

Description	mTOR inhibitor-8 is a potent mTOR inhibitor and autophagy inducer with antiviral and antitumor activity. mTOR inhibitor-8 inhibits the growth of A549 cells, which can be used to study non-small cell lung cancer.
Targets(IC50)	Autophagy,mTOR
In vitro	mTOR inhibitor-8 (Compound 5e; 0.1-10µM) suppresses the growth of A549 cells in a dose-dependent manner after 24 and 48 hours of treatment[1]. At a concentration of 10 µM, mTOR inhibitor-8 induces autophagy in an mTOR-dependent and time-dependent manner (3-24hours), increasing LC3B-II levels and reducing the phosphorylation of RPS6KB1 (ribosomal protein S6 kinase) and EIF4EBP1 (eukaryotic translation initiation factor 4E-binding protein 1)[1].
In vivo	In vivo, mTOR inhibitor-8 effectively inhibits tumor growth at concentrations of 25 and 50 µM for a duration of 6 days, without adversely affecting angiogenesis in the normal chick chorioallantoic membrane [1].

Solubility Information

Solubility	DMSO: 20 mg/mL (44.75 mM),Sonication is recommended. (< 1 mg/ml refers to the product slightly soluble or insoluble)
In vivo Formulation	10% DMSO+90% Saline: 2 mg/mL (4.47 mM),Sonication is recommended. <i>Please add the solvents sequentially, clarifying the solution as much as possible before adding the next one. Dissolve by heating and/or sonication if necessary. Working solution is recommended to be prepared and used immediately. The formulation provided above is for reference purposes only. In vivo formulations may vary and should be modified based on specific experimental conditions.</i>

Preparing Stock Solutions

	1mg	5mg	10mg
1 mM	2.2374 mL	11.1869 mL	22.3739 mL
5 mM	0.4475 mL	2.2374 mL	4.4748 mL
10 mM	0.2237 mL	1.1187 mL	2.2374 mL
50 mM	0.0447 mL	0.2237 mL	0.4475 mL

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. Please use it as soon as possible.

Note: The dilution table applies only to solid products. For liquid products, please calculate the stock solution based on the stated concentration and/or density.

Reference

ZhaoMin Lin, et al. Discovery of new fluorescent thiazole-pyrazoline derivatives as autophagy inducers by inhibiting mTOR activity in A549 human lung cancer cells. Cell Death Dis. 2020 Jul 20;11(7):551.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481