


Wiskostatin

Chemical Properties

CAS No. :	253449-04-6
Formula:	C ₁₇ H ₁₈ Br ₂ N ₂ O
Molecular Weight:	426.15
Storage:	Powder: -20°C for 3 years In solvent: -80°C for 1 year Actual storage temperature shall be subject to the COA.



Biological Description

Description	Wiskostatin is a selective inhibitor of N-WASP, a ubiquitously expressed member of the Wiskott-Aldrich Syndrome protein (WASp) family that regulates actin polymerization.
Targets(IC50)	Others, Arp2/3 Complex
Kinase Assay	Enzyme kinetic analysis is performed on the purified wild type and mutant UL97 protein species using increasing concentrations of ATP (2 μ M to 20 μ M). The amount of incorporated radiolabelled phosphate is plotted against the concentration of ATP in a Lineweaver Burke plot to determine the Km for ATP for each UL97 species. The effect of Maribavir upon the rate of radiolabelled phosphate incorporation by wild type or mutant UL97 is determined by protein kinase assays at a fixed concentration of Maribavir (0.5 μ M) as above, or with increasing concentrations of Maribavir (0.01 μ M to 5.0 μ M) to determine the IC50 of Maribavir for each UL97 species. In order to determine the nature of the inhibition mediated by Maribavir, plots of 1/v vs 1/ATP with increasing concentrations of Maribavir are constructed. Competitive inhibition is evident if the family of lines converged on the y-axis at 1/Vmax. The change in slope caused by the addition of Maribavir is used to calculate the Ki[1].

Solubility Information

Solubility	DMSO: 76.9 mg/mL (180.45 mM), Sonication is recommended. (< 1 mg/ml refers to the product slightly soluble or insoluble)
In vivo Formulation	10% DMSO+90% Saline: < 7.69 mg/mL (18.05 mM), Lower concentrations may be soluble, but exact solubility limit is unknown. 10% DMSO+90% Corn Oil: 2.5 mg/mL (5.87 mM), Sonication is recommended. 10% DMSO+40% PEG300+5% Tween 80+45% Saline: 7.69 mg/mL (18.05 mM), Solution. <i>Please add the solvents sequentially, clarifying the solution as much as possible before adding the next one. Dissolve by heating and/or sonication if necessary. Working solution is recommended to be prepared and used immediately. The formulation provided above is for reference purposes only. In vivo formulations may vary and should be modified based on specific experimental conditions.</i>

Preparing Stock Solutions

	1mg	5mg	10mg
1 mM	2.3466 mL	11.733 mL	23.4659 mL
5 mM	0.4693 mL	2.3466 mL	4.6932 mL
10 mM	0.2347 mL	1.1733 mL	2.3466 mL
50 mM	0.0469 mL	0.2347 mL	0.4693 mL

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. Please use it as soon as possible.

Note: The dilution table applies only to solid products. For liquid products, please calculate the stock solution based on the stated concentration and/or density.

Reference

Guerrero CJ, Weisz OA. Am J Physiol Cell Physiol. 2007 Apr;292(4):C1562-6.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481