

## Glycitein

## Chemical Properties

CAS No. : 40957-83-3


Formula: C<sub>16</sub>H<sub>12</sub>O<sub>5</sub>

Molecular Weight: 284.26

Storage: Keep away from direct sunlight, Store at low temperature

Powder: -20°C for 3 years | In solvent: -80°C for 1 year

Actual storage temperature shall be subject to the COA.



## Biological Description

Description	1. Glycitein has weak estrogenic activity. 2. Glycitein (Glycetein) inhibits glioma cell invasion through down-regulation of MMP-3 and MMP-9 gene expression.
Targets(IC50)	Apoptosis, Autophagy

## Solubility Information

Solubility	DMSO: 25.75 mg/mL (90.59 mM), Sonication is recommended. (< 1 mg/ml refers to the product slightly soluble or insoluble)
In vivo Formulation	10% DMSO+90% Corn Oil: 2.5 mg/mL (8.79 mM), Sonication is recommended. <i>Please add the solvents sequentially, clarifying the solution as much as possible before adding the next one. Dissolve by heating and/or sonication if necessary. Working solution is recommended to be prepared and used immediately. The formulation provided above is for reference purposes only. In vivo formulations may vary and should be modified based on specific experimental conditions.</i>

## Preparing Stock Solutions

	1mg	5mg	10mg
1 mM	3.5179 mL	17.5895 mL	35.1791 mL
5 mM	0.7036 mL	3.5179 mL	7.0358 mL
10 mM	0.3518 mL	1.759 mL	3.5179 mL
50 mM	0.0704 mL	0.3518 mL	0.7036 mL

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. Please use it as soon as possible.

Note: The dilution table applies only to solid products. For liquid products, please calculate the stock solution based on the stated concentration and/or density.

### Reference

Pan W , Ikeda K , Takebe M , et al. Genistein, daidzein and glycitein inhibit growth and DNA synthesis of aortic smooth muscle cells from stroke-prone spontaneously hypertensive rats.[J]. Journal of Nutrition, 2001, 131(4): 1154-8.

Ferreira R S B, Farias F O, de Araujo E J S, et al. Deep eutectic solvents as an alternative for extraction of flavonoids from soybean (Glycine max (L) Merrill) and okara: An experimental and computational approach based on COSMO-SAC model. Food Research International. 2023: 113266.

**Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins**

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel: 781-999-4286 E\_mail: info@targetmol.com Address: 34 Washington Street, Wellesley Hills, MA 02481