

Thymine

Chemical Properties


CAS No. : 533-67-5

Formula: C₅H₁₀O₄

Molecular Weight: 134.13

Storage: Powder: -20°C for 3 years | In solvent: -80°C for 1 year

Actual storage temperature shall be subject to the COA.



Biological Description

Description	Thymine induces apoptosis by inhibiting glutathione synthesis and increasing its efflux.
Targets(IC50)	Apoptosis,Endogenous Metabolite

Solubility Information

Solubility	DMSO: 55 mg/mL (410.05 mM),Sonication is recommended. (< 1 mg/ml refers to the product slightly soluble or insoluble)
In vivo Formulation	10% DMSO+40% PEG300+5% Tween 80+45% Saline: 2 mg/mL (14.91 mM),Sonication is recommended. <i>Please add the solvents sequentially, clarifying the solution as much as possible before adding the next one. Dissolve by heating and/or sonication if necessary. Working solution is recommended to be prepared and used immediately. The formulation provided above is for reference purposes only. In vivo formulations may vary and should be modified based on specific experimental conditions.</i>

Preparing Stock Solutions

	1mg	5mg	10mg
1 mM	7.4555 mL	37.2773 mL	74.5545 mL
5 mM	1.4911 mL	7.4555 mL	14.9109 mL
10 mM	0.7455 mL	3.7277 mL	7.4555 mL
50 mM	0.1491 mL	0.7455 mL	1.4911 mL

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. Please use it as soon as possible.

Note: The dilution table applies only to solid products. For liquid products, please calculate the stock solution based on the stated concentration and/or density.

Reference

Fico A, et al. 2-deoxy-d-ribose induces apoptosis by inhibiting the synthesis and increasing the efflux of glutathione. Free Radic Biol Med. 2008 Jul 15;45(2):211-7.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481