

CRT0044876

Chemical Properties


CAS No. : 6960-45-8

Formula: C₉H₆N₂O₄

Molecular Weight: 206.15

Storage: Powder: -20°C for 3 years | In solvent: -80°C for 1 year

Actual storage temperature shall be subject to the COA.



Biological Description

Description	CRT0044876 (7-NO ₂ -ICA) is a potent and selective APE1 inhibitor with IC ₅₀ of ~3 μM.
Targets(IC ₅₀)	DNA/RNA Synthesis
In vitro	CRT0044876 potently inhibits the AP endonuclease, 3'-phosphodiesterase and 3'-phosphatase activities of APE1, and exhibits the selectivity for the inhibition of exonuclease III family. CRT0044876 potentiates the cytotoxicity of several DNA base-targeting compounds with an accumulation of unrepaired AP sites. [1] CRT0044876, by inhibition of the BER pathway increases oxidative DNA damage temporally related to increased intracellular reactive oxygen species in an acidic tumor microenvironment, and thus results in cell cycle arrests and increased DNA double strand breaks, leading to cell death. [2]
Kinase Assay	AP site cleavage assay: BER reaction buffer comprises 40 mM HEPES-KOH (pH 7.8), 5 mM MgCl ₂ , 0.5 mM DTT and 0.1 mM EDTA. A 10 μl AP site cleavage reaction comprised of BER buffer mix, purified protein (3.3 nM final concentration of APE1) and 0.75 ng reduced AP site double-stranded oligonucleotide. The mixture is incubated at 37°C for 1 h. A total of 1 μl of stop buffer (50% glycerol, 10 mM Tris-HCl, 1 mM EDTA, 0.1% bromophenol blue and 0.1% Xylene cyanol) is added, and the sample mixture is denatured at 90-100°C for 2 min. The sample is then run on a 15% TBE Criterion? Pre-Cast Gel, with electrophoresis at a constant current of 30 mA for 30 min, and the radiolabeled substrate and reaction products are visualized using a phosphorImager. The inhibitory activity of potential APE1-targeting compounds are analyzed at drug concentrations ranging from 0.1 to 100 μM. The resolved radiolabeled bands are quantified using ImageQuant software analysis, and IC ₅₀ values are calculated by determining the concentration of the inhibitor that reduced APE1 activity to 50% of the control values.
Cell Research	HT1080 fibrosarcoma cells are grown in 2% RPMI medium [supplemented with penicillin 0.06 g/l, streptomycin 0.1 g/l (pH 7.0), 10% fetal bovine serum and 4 mM glutamine]. Only cells with a plating efficiency of ≥60% are used for clonogenic survival analysis. Tissue culture dishes (10 cm) are seeded with 500 cells per dish and the culture is maintained in a humidified incubator at 37°C in an atmosphere of 5% CO ₂ and 95% air. To evaluate the toxicity profile of putative APE1 inhibitors, various concentrations (100-500 μM) of inhibitor are added to the medium, and cultures were incubated for 7-10 days until cell colonies are formed. Colonies are fixed [75% (v/v) methanol, 25% (v/v)

A DRUG SCREENING EXPERT

Cell Research	acetic acid] for 30 min and stained with crystal violet (1 mg/ml in distilled water) for 4 h at room temperature. Visible colonies are counted on a colony counter. (Only for Reference)
---------------	--

Solubility Information

Solubility	DMSO: 100 mg/mL (485.08 mM),Sonication is recommended. Ethanol: 1 mg/mL (4.85 mM),Sonication is recommended. H2O: < 1 mg/mL (insoluble or slightly soluble), (< 1 mg/ml refers to the product slightly soluble or insoluble)
In vivo Formulation	10% DMSO+40% PEG300+5% Tween 80+45% Saline: 2 mg/mL (9.7 mM),Sonication is recommended. <i>Please add the solvents sequentially, clarifying the solution as much as possible before adding the next one. Dissolve by heating and/or sonication if necessary. Working solution is recommended to be prepared and used immediately. The formulation provided above is for reference purposes only. In vivo formulations may vary and should be modified based on specific experimental conditions.</i>

Preparing Stock Solutions

	1mg	5mg	10mg
1 mM	4.8508 mL	24.2542 mL	48.5084 mL
5 mM	0.9702 mL	4.8508 mL	9.7017 mL
10 mM	0.4851 mL	2.4254 mL	4.8508 mL
50 mM	0.097 mL	0.4851 mL	0.9702 mL

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. Please use it as soon as possible.

Note: The dilution table applies only to solid products. For liquid products, please calculate the stock solution based on the stated concentration and/or density.

Reference

Madhusudan S, et al. Nucleic Acids Res. 2005, 33(15), 4711-4724.

Long K, Gu L, Li L, et al. Small-molecule inhibition of APE1 induces apoptosis, pyroptosis, and necroptosis in non-small cell lung cancer. Cell Death & Disease. 2021, 12(6): 1-15.

Du Y, Zhou Y, Yan X, et al. APE1 inhibition enhances ferroptotic cell death and contributes to hepatocellular carcinoma therapy. Cell Death & Differentiation. 2024: 1-16.

Seo Y, et al. Cancer Res. 2009, 69(18), 7285-7293.

Long K, Gu L, Li L, et al. Small-molecule inhibition of APE1 induces apoptosis, pyroptosis, and necroptosis in non-small cell lung cancer[J]. Cell Death & Disease. 2021, 12(6): 1-15.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel: 781-999-4286 E_mail: info@targetmol.com Address: 34 Washington Street, Wellesley Hills, MA 02481