

## Cyclo(Gly-L-Pro)


## Chemical Properties

CAS No. : 3705-27-9

Formula: C7H10N2O2

Molecular Weight: 154.17

Storage: Keep away from moisture, Store at low temperature,  
Keep away from direct sunlight  
Powder: -20°C for 3 years | In solvent: -80°C for 1 year  
Actual storage temperature shall be subject to the COA.



## Biological Description

Description	Cyclo(Gly-L-Pro), a cyclic dipeptide, demonstrates significant immunomodulatory activity by stimulating beneficial immune responses while simultaneously suppressing the release of the pro-inflammatory cytokine TNF- $\alpha$ , reducing mRNA expression of IL-1 $\beta$ and IL-6, and inhibiting nitric oxide (NO) production. It has also been shown to enhance memory and improve functional recovery in animal models, indicating substantial potential for neuroscience research due to its neuroprotective and immunostimulatory properties.
Targets(IC50)	TNF
In vitro	<b>Methods:</b> CIK cells were stimulated with Cyclo(Gly-L-Pro) (1, 10, 100 $\mu$ g/ml), and the expression of MyD88, IL-1 $\beta$ , TNF- $\alpha$ , I-IFN, and IL-8 genes at different time points (2, 8, and 24 h) after stimulation was quantified by real-time fluorescence. Lipopolysaccharide (LPS), a known immunostimulant, was used as a positive control. To analyze whether these compounds were toxic to the cells, changes in cell viability were measured using a methyltetrazole assay. <b>Results:</b> Cyclo(Gly-L-Pro) and LPS significantly increased the transcription level of MyD88 as early as 2 h after exposure; Cyclo(Gly-L-Pro) also induced the gene expression of cytokines such as IL-1 $\beta$ , TNF- $\alpha$ , and I-IFN. Under the experimental conditions, Cyclo(Gly-L-Pro) was not toxic to CIK cells. [1]

## Solubility Information

Solubility	H2O: 180 mg/mL (1167.54 mM), Sonication is recommended. DMSO: 20 mg/mL (129.73 mM), Sonication is recommended. ( $< 1$ mg/ml refers to the product slightly soluble or insoluble)
------------	---

### Preparing Stock Solutions

---

	1mg	5mg	10mg
1 mM	6.4863 mL	32.4317 mL	64.8635 mL
5 mM	1.2973 mL	6.4863 mL	12.9727 mL
10 mM	0.6486 mL	3.2432 mL	6.4863 mL
50 mM	0.1297 mL	0.6486 mL	1.2973 mL

---

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. Please use it as soon as possible.

Note: The dilution table applies only to solid products. For liquid products, please calculate the stock solution based on the stated concentration and/or density.

### Reference

Wu ZF, et al. Induction of immune-related gene expression in *Ctenopharyngodon idella* kidney cells by secondary metabolites from immunostimulatory *Alcaligenes faecalis* FY-3. *Scand J Immunol.* 2012 Aug;76(2):131-40.

Khan R, et al. Attenuation of TNF- $\alpha$  secretion by L-proline-based cyclic dipeptides produced by culture broth of *Pseudomonas aeruginosa*. *Bioorg Med Chem Lett.* 2015 Dec 15;25(24):5756-61.

**Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins**

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E\_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481