


Nabumetone

Chemical Properties

CAS No. :	42924-53-8
Formula:	C ₁₅ H ₁₆ O ₂
Molecular Weight:	228.29
Storage:	Powder: -20°C for 3 years In solvent: -80°C for 1 year Actual storage temperature shall be subject to the COA.



Biological Description

Description	Nabumetone (BRL14777)(BRL14777) is a non-steroidal anti-inflammatory drug and its active metabolite 6MNA inhibits the enzymes cyclo-oxygenase I and II.
Targets(IC50)	Glutathione Peroxidase, COX
In vitro	Nabumetone exhibits anti-inflammatory activity in rat models of carrageenan-induced edema and UV-induced erythema in guinea pigs. In a rat granuloma model, Nabumetone is non-toxic at concentrations higher than the minimum effective dose. Furthermore, in adjuvant-induced arthritic rats, Nabumetone demonstrates good gastric tolerability at doses exceeding those required for anti-inflammatory efficacy.
In vivo	Nabumetone undergoes extensive first-pass metabolism to form its main active metabolite (6-MNA) in the systemic circulation, which is capable of inhibiting COX-2.
Kinase Assay	Cells are harvested, washed, and lysed in NP-40 lysis buffer (50 mM Tris-HCl [pH 7.5], 150 mM NaCl, 0.5% NP-40). Total cell protein is quantified using the Bradford assay and 1-mg/mL protein aliquots are made. Ten microliters of total cell protein is mixed with 290 µL of substrate solution (0.1 mg/mL dithiothreitol [DTT], 0.1 mg/mL albumin, and 1 mM alanine-β-naphthylamide). Fluorometric measurements (340 nm excitation, 400 nm emission) are made after 15 and 30 min. The slope of the line between the 15- and 30-min measurements is used to represent aminopeptidase activity. Total cell protein is preincubated with bestatin, amastatin, puromycin, EDTA, and/or ZnCl ₂ for 20 min before the fluorometric aminopeptidase assay.

Solubility Information

Solubility	DMSO: 55 mg/mL (240.92 mM), Sonication is recommended. Ethanol: 24 mg/mL (105.13 mM), Sonication is recommended. (< 1 mg/ml refers to the product slightly soluble or insoluble)
In vivo Formulation	10% DMSO+40% PEG300+5% Tween 80+45% Saline: 2 mg/mL (8.76 mM), Sonication is recommended. <i>Please add the solvents sequentially, clarifying the solution as much as possible before adding the next one. Dissolve by heating and/or sonication if necessary. Working solution is recommended to be prepared and used immediately. The formulation provided above is for reference purposes only. In vivo formulations may vary and should be modified based on specific experimental conditions.</i>

Preparing Stock Solutions

	1mg	5mg	10mg
1 mM	4.3804 mL	21.902 mL	43.8039 mL
5 mM	0.8761 mL	4.3804 mL	8.7608 mL
10 mM	0.438 mL	2.1902 mL	4.3804 mL
50 mM	0.0876 mL	0.438 mL	0.8761 mL

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. Please use it as soon as possible.

Note: The dilution table applies only to solid products. For liquid products, please calculate the stock solution based on the stated concentration and/or density.

Reference

Hedner T, et al. *Drugs*, 2004, 64(20), 2315-2343.

Boyle EA, et al. *J Pharm Pharmacol*, 1982, 34(9), 562-569.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481