


AZD-1480

Chemical Properties

CAS No. :	935666-88-9
Formula:	C ₁₄ H ₁₄ ClFN ₈
Molecular Weight:	348.77
Storage:	Powder: -20°C for 3 years In solvent: -80°C for 1 year Actual storage temperature shall be subject to the COA.



Biological Description

Description	AZD1480 is a novel ATP-competitive JAK2 inhibitor (IC ₅₀ : 0.26 nM), selectivity act against Tyk2 and JAK3, and to a smaller extent against JAK1. AZD1480 has been used in trials studying the treatment of Solid Malignancies, Post-Polycythaemia Vera, Primary Myelofibrosis (PMF), and Essential Thrombocythaemia Myelofibrosis.
Targets(IC ₅₀)	JAK
In vitro	AZD1480 inhibits the growth of subcutaneous xenograft tumors and extends the lifespan of mice carrying intracranial malignant gliomas by inhibiting the activity of STAT3, indicating that AZD1480's inhibition of the JAK/STAT3 pathway can be utilized in research targeting the treatment of patients with intracranial malignant gliomas. Furthermore, AZD1480 suppresses STAT3 phosphorylation in human solid tumor and multiple myeloma models, reduces angiogenesis and metabolism in human xenograft models, and demonstrates inhibitory effects on the growth of human solid tumors with active STAT. Additionally, AZD1480 inhibits lung metastasis and pulmonary infiltration of tumors in bone marrow cells.
In vivo	AZD1480, by affecting the microenvironment of tumors, can inhibit local metastasis and angiogenesis. As an effective inhibitor of JAK2, it suppresses the growth and survival of human multiple myeloma cells, as well as inhibits FGFR3 and STAT3 signaling and downstream targets, including Cyclin D2. At low micromolar concentrations, AZD1480 induces apoptosis and inhibits cell proliferation in myeloma cell lines. Moreover, at 5 μM, it induces cell cycle arrest at the G ₂ /M phase and cell death through inhibition of Aurora kinases. In both human and murine glioma cells, AZD1480 effectively inhibits constitutive and inducible phosphorylation of JAK1/2 and STAT-3, thereby reducing cell proliferation and inducing apoptosis.
Kinase Assay	kinase assays: Inhibition studies of AZD1480 are performed using recombinant Jak1, Jak2, or Jak3 under buffer conditions of 50 mM HEPES pH 7.3, 1 mM DTT, 0.01% Tween-20, 50 mM/ml BSA, and 10 mM MgCl ₂ . Jak3 enzyme is expressed as N-terminal GST fusion in insect cells and purified by glutathione-affinity and size-exclusion chromatographies. Enzymes are assayed in the presence of AZD1480 (10 point dose response, in triplicate, from 8.3 μM to 0.3 nM in half-log dilution steps) using 1.5 μM peptide substrate (Jak1: FITC-C6-KKHTDDGYMPMSPGVA-NH ₂ , Jak2 and Jak3: FAM-SRCtide) and screened under their respective ATP Km (Jak1: 55 μM, Jak2: 15 μM, Jak3: 3 μM) and approximated physiological ATP concentration of 5 mM. Phosphorylated and unphosphorylated peptides are separated and quantified by a Caliper LC3000 system

Kinase Assay	for calculating percent inhibition.
Cell Research	Renca or 786-O cells are suspended in DMEM medium with 5% FBS , and seeded in 96-well plates (5×10 ³ per well) to allow adhesion and then treated with DMSO or AZD1480 for 48 hours. Cell viability is determined by MTS assay. Absorbance at 490 nm is measured with Mikrotek Laborsysteme. Mouse endothelial cells and splenic CD11b+/c-myeloid cells are enriched from tumor-bearing mice, and cultured in 5% FBS RPMI-1640 medium. HUVECs are cultured on collagen 1-coated plates in complete medium. All cells are treated with DMSO and AZD1480 at various doses for 24 hours. Cell viability is determined by counting cell number manually. All the experiments are repeated 3 times. (Only for Reference)

Solubility Information

Solubility	Ethanol: < 1 mg/mL (insoluble or slightly soluble), H ₂ O: < 1 mg/mL (insoluble or slightly soluble), DMSO: 72.5 mg/mL (207.87 mM), Sonication is recommended. (< 1 mg/ml refers to the product slightly soluble or insoluble)
In vivo Formulation	10% DMSO+90% Saline: < 7.25 mg/mL (20.79 mM), Lower concentrations may be soluble, but exact solubility limit is unknown. 10% DMSO+40% PEG300+5% Tween 80+45% Saline: 7.25 mg/mL (20.79 mM), Solution. <i>Please add the solvents sequentially, clarifying the solution as much as possible before adding the next one. Dissolve by heating and/or sonication if necessary. Working solution is recommended to be prepared and used immediately. The formulation provided above is for reference purposes only. In vivo formulations may vary and should be modified based on specific experimental conditions.</i>

Preparing Stock Solutions

	1mg	5mg	10mg
1 mM	2.8672 mL	14.3361 mL	28.6722 mL
5 mM	0.5734 mL	2.8672 mL	5.7344 mL
10 mM	0.2867 mL	1.4336 mL	2.8672 mL
50 mM	0.0573 mL	0.2867 mL	0.5734 mL

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. Please use it as soon as possible.

Note: The dilution table applies only to solid products. For liquid products, please calculate the stock solution based on the stated concentration and/or density.

Reference

E Derenzini, et al. *Blood Cancer J*, 2011, 1(12), e46.

Yang T, Shi X, Kang Y, et al. Novel compounds TAD-1822-7-F2 and F5 inhibited HeLa cells growth through the JAK/Stat signaling pathway. *Biochemical & Pharmacology*. 2018, 103: 118-126

Hu Z, Sui Q, Jin X, et al. IL6-STAT3-C/EBP β -IL6 positive feedback loop in tumor-associated macrophages promotes the EMT and metastasis of lung adenocarcinoma. *Journal of Experimental & Clinical Cancer Research*. 2024, 43(1): 63.

Scuto A, et al. *Leukemia*, 2011, 25(3), 538-50.

McFarland BC, et al. *Cancer Ther*, 2011, 10(12), 2384-93.

Xin H, et al. *Cancer Res*, 2011, 71(21), 6601-10.

Hedvat M, et al. *Cancer Cell*, 2009, 16(6), 487-97.

Ni J, et al. Tyrosine receptor kinase B is a drug target in astrocytomas. *Neuro Oncol*. 2017 Jan;19(1):22-30.

Yang, Tianfeng, Xianpeng Shi, Yuan Kang, Man Zhu, Mengying Fan, Dongdong Zhang, and Yanmin Zhang. Novel compounds TAD-1822-7-F2 and F5 inhibited HeLa cells growth through the JAK/Stat signaling pathway [J].

Biomedicine & Pharmacotherapy . 2018 Jul; 103:118-126.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel: 781-999-4286

E_mail: info@targetmol.com

Address: 34 Washington Street, Wellesley Hills, MA 02481