


## Pelitinib

## Chemical Properties

CAS No. :	257933-82-7
Formula:	C <sub>24</sub> H <sub>23</sub> ClFN <sub>5</sub> O <sub>2</sub>
Molecular Weight:	467.92
Storage:	Powder: -20°C for 3 years   In solvent: -80°C for 1 year Actual storage temperature shall be subject to the COA.



## Biological Description

Description	Pelitinib (EKB-569) (EKB-569) is an effective irreversible EGFR inhibitor (IC <sub>50</sub> : 38.5 nM).
Targets(IC <sub>50</sub> )	EGFR,Raf,MEK,Src
In vitro	In the A431 xenograft model targeting EGFR, Pelitinib (10 mg/kg, p.o.) effectively inhibits EGFR phosphorylation, achieving a 90% inhibition within 1 hour and maintaining over 50% inhibition after 24 hours. In the mouse airway epithelial cell remodeling model, which can be induced by viral infection and is characterized by a delayed yet permanent transition to goblet cell metaplasia, Pelitinib (20 mg/kg/day) completely blocks the increase of ciliated cells and the decrease of Clara cells, thereby correcting epithelial cell remodeling from three different aspects and significantly inhibiting the occurrence of goblet cells. Additionally, Pelitinib (20 mg/kg/day) inhibits 87% of tumor occurrence in APCMin/+ mice, which is comparable to the effect of a double dose of EKI-785 (40 mg/kg/day), consistent with its higher in vivo activity. Pelitinib exhibits selective inhibitory effects on EGFR signaling in airway epithelial cells in vivo.
In vivo	Pelitinib (75-500 nM) specifically inhibits the activation of AKT and ERK1/2 without affecting the NF-κB pathway. In normal human keratinocytes, Pelitinib effectively inhibits TGF-α-mediated EGFR activation (IC <sub>50</sub> : 56 nM) and the activation of STAT3 and ERK1/2 (IC <sub>50</sub> : 60 nM and 62 nM, respectively). It also suppresses EGF-induced EGFR phosphorylation (IC <sub>50</sub> : 20-80 nM) and STAT3 phosphorylation (IC <sub>50</sub> : 30-70 nM) in both A431 cells and normal human keratinocytes. Moreover, Pelitinib significantly inhibits the proliferation of normal human keratinocytes (IC <sub>50</sub> : 61 nM), A431 (IC <sub>50</sub> : 125 nM), and MDA-468 (IC <sub>50</sub> : 260 nM) tumor cells, but shows lower activity against MCF-7 cells (IC <sub>50</sub> : 3.6 μM).
Kinase Assay	Autophosphorylation of EGFR in cells: For experiments using cells in culture, A431 cells are treated with various concentrations of Pelitinib for 2.75 hours before co-incubation with 100 ng/mL EGF for 0.25 hour. Cells are washed twice with cold phosphate-buffered saline (PBS) before adding to lysis buffer (10 mM Tris, pH 7.5, 5 mM ethylenediamine tetra-acetic acid (EDTA), 150 mM NaCl, 1% Triton X-100, 1% Sodium deoxycholate, 0.1 % SDS, 1 mM PMSF, 10 mg/mL pepstatin A, 10 mg/mL leupeptin, 20 KIU/mL aprotinin, 2 mM sodium orthovanadate, and 100 mM sodium fluoride) for 20 minutes on ice, before immunoprecipitation and SDS-PAGE-immunoblotting. For immunoprecipitation, cultured cells are placed in cold lysis buffer and immediately homogenized on ice with a

Kinase Assay	polytron with several pulses. The homogenate is first centrifuged at 2500 rpm (20 minutes, 4 °C) and then again at 14,000 rpm in a microcentrifuge (10 minutes, 4 °C). Supernatants (1000 µg protein) are incubated for 2 hours at 4 °C with 15 mL of EGFR polyclonal antibody. After 2 hours, 50 µL of protein G plus/protein A agarose beads is added and incubated with constant rotation for 2 hours at 4 °C. After washing with lysis buffer, beads are boiled for 2 minutes in Laemmli sample buffer. Proteins are then resolved by SDS-PAGE, transferred to immobilon membrane and probed overnight with an anti-phosphotyrosine antibody conjugated with horseradish peroxidase (HRP). Membranes are developed using the ECL reagent. Total EGFR protein is determined by stripping membranes and re-probing with receptor-specific antibodies. Quantitation of bands is done by densitometry, using ImageQuant software with a Molecular Dynamics laser transmittance scanner.
Cell Research	Cells are seeded in 96-well dishes, and after 2 hours, Pelitinib is added and incubated for 5 days. After incubation, the medium is removed from each well and fresh medium (150 µL) + 1 mg/mL MTT solution (50 µL) is added. After incubation for 2 hours at 37 °C, the medium is replaced with 150 µL DMSO, and absorbance at 540 nm in each well is determined. The IC50 is calculated by linear regression of the data. (Only for Reference)

### Solubility Information

Solubility	H2O: < 1 mg/mL (insoluble or slightly soluble), Ethanol: < 1 mg/mL (insoluble or slightly soluble), DMSO: 9.43 mg/mL (20.15 mM),Sonication is recommended. (< 1 mg/ml refers to the product slightly soluble or insoluble)
In vivo Formulation	10% DMSO+40% PEG300+5% Tween 80+45% Saline: 1 mg/mL (2.14 mM),Sonication is recommended. <i>Please add the solvents sequentially, clarifying the solution as much as possible before adding the next one. Dissolve by heating and/or sonication if necessary. Working solution is recommended to be prepared and used immediately. The formulation provided above is for reference purposes only. In vivo formulations may vary and should be modified based on specific experimental conditions.</i>

### Preparing Stock Solutions

	1mg	5mg	10mg
1 mM	2.1371 mL	10.6856 mL	21.3712 mL
5 mM	0.4274 mL	2.1371 mL	4.2742 mL
10 mM	0.2137 mL	1.0686 mL	2.1371 mL
50 mM	0.0427 mL	0.2137 mL	0.4274 mL

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. Please use it as soon as possible.

Note: The dilution table applies only to solid products. For liquid products, please calculate the stock solution based on the stated concentration and/or density.

Reference

Torrance CJ, et al. Nat Med, 2000, 6(9), 1024-1028.

Li J, Zheng M, Xu Y, et al. Target Proteins Profiling of Irreversible Kinase Inhibitor Pelitinib and Discovery of Degradation of PRDX4 by Label Free Chemoproteomics. Journal of Pharmaceutical and Biomedical Analysis. 2023: 115398.

Nunes M, et al. Mol Cancer Ther, 2004, 3(1), 21-27.

Tyner JW, et al. J Clin Invest, 2006, 116(2), 309-321.

**Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins**

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel: 781-999-4286 E\_mail: info@targetmol.com Address: 34 Washington Street, Wellesley Hills, MA 02481