


## Doxycycline

## Chemical Properties

CAS No. :	564-25-0
Formula:	C <sub>22</sub> H <sub>24</sub> N <sub>2</sub> O <sub>8</sub>
Molecular Weight:	444.43
Storage:	Powder: -20°C for 3 years   In solvent: -80°C for 1 year Actual storage temperature shall be subject to the COA.



## Biological Description

Description	Doxycycline is an orally active tetracycline antibiotic with broad-spectrum inhibitory effects on matrix metalloproteinases (MMPs), possessing both antibacterial and antitumor activities, and is commonly used in inducible gene expression ON-OFF systems.
Targets(IC50)	MMP, ribosome, Antibacterial, Antibiotic, Parasite
In vitro	<p><b>METHODS:</b> Twenty-two tumor cells were treated with Doxycycline for 48 h. Cell viability was measured by MTT.</p> <p><b>RESULTS:</b> Lung cancer cells were more sensitive to Doxycycline than most other cell lines. NCI-H446 and A549 cells were sensitive to Doxycycline with IC50s of 1.70 <math>\mu</math>M and 1.06 <math>\mu</math>M, respectively. [1]</p> <p><b>METHODS:</b> Human pancreatic cancer cells PANC-1 were treated with Doxycycline (10-40 <math>\mu</math>g/mL) for 8-24 h. The cell cycle was detected by Flow Cytometry.</p> <p><b>RESULTS:</b> Doxycycline caused the cells to arrest in the G1-S phase of the cell cycle. [2]</p>
In vivo	<p><b>METHODS:</b> To detect anti-tumor activity in vivo, Doxycycline (50 mg/kg) was administered intraperitoneally once daily for fifteen days to BALB/c mice harboring human mammary carcinoma tumor 4T1.</p> <p><b>RESULTS:</b> Doxycycline delayed tumor growth and reduced the number of tumor-associated macrophages and blood vessels. [3]</p> <p><b>METHODS:</b> To assay antimicrobial activity in vivo, Doxycycline (40 mg/kg) was administered intraperitoneally to BALB/c mice exposed to <i>B. pseudomallei</i> 1026b twice daily for two weeks.</p> <p><b>RESULTS:</b> Doxycycline-treated mice survived 100% of the time, and no <i>B. pseudomallei</i> DNA was amplified from the lungs or spleens of the majority of the surviving mice. [4]</p>

## Solubility Information

Solubility	H <sub>2</sub> O: < 1 mg/mL (insoluble) DMSO: 200 mg/mL (450.01 mM), Sonication is recommended. (< 1 mg/ml refers to the product slightly soluble or insoluble)
In vivo Formulation	10% DMSO+90% Saline: 4 mg/mL (9 mM), Solution. <i>Please add the solvents sequentially, clarifying the solution as much as possible before adding the next one. Dissolve by heating and/or sonication if necessary. Working solution is recommended to be prepared and used immediately. The formulation provided above is for reference purposes only. In vivo formulations may</i>

## A DRUG SCREENING EXPERT

In vivo Formulation	<i>vary and should be modified based on specific experimental conditions.</i>
---------------------	---

### Preparing Stock Solutions

	1mg	5mg	10mg
1 mM	2.2501 mL	11.2504 mL	22.5007 mL
5 mM	0.450 mL	2.2501 mL	4.5001 mL
10 mM	0.225 mL	1.125 mL	2.2501 mL
50 mM	0.045 mL	0.225 mL	0.450 mL

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. Please use it as soon as possible.

Note: The dilution table applies only to solid products. For liquid products, please calculate the stock solution based on the stated concentration and/or density.

### Reference

- Qin Y, et al. Doxycycline reverses epithelial-to-mesenchymal transition and suppresses the proliferation and metastasis of lung cancer cells. *Oncotarget*. 2015 Dec 1;6(38):40667-79.
- Tan Y, Chen X, Xu Z, et al. The mechanics of tumor cells dictate malignancy via cytoskeleton-mediated APC/Wnt/ $\beta$ -catenin signaling. *Research*. 2023
- Wang J, Zhang B, Chen X, et al. Cell mechanics regulate the migration and invasion of hepatocellular carcinoma cells via JNK signaling. *Acta Biomaterialia*. 2024
- Li Y M, Mei Y C, Liu A H, et al. Gcn5- and Bre1-mediated Set2 degradation promotes chronological aging of *Saccharomyces cerevisiae*. *Cell Reports*. 2023, 42(10).
- Son K, et al. Doxycycline induces apoptosis in PANC-1 pancreatic cancer cells. *Anticancer Res*. 2009 Oct;29(10):3995-4003.
- Tang X, et al. Doxycycline attenuates breast cancer related inflammation by decreasing plasma lysophosphatidate concentrations and inhibiting NF- $\kappa$ B activation. *Mol Cancer*. 2017 Feb 8;16(1):36.
- Wang W, Wang M, Du T, et al. SHMT2 Promotes Gastric Cancer Development through Regulation of HIF1 $\alpha$ /VEGF/STAT3 Signaling. *International Journal of Molecular Sciences*. 2023, 24(8): 7150.
- Lu J, Lu Z, Liu L, et al. Identification of crocin as a new hIAPP amyloid inhibitor via a simple yet highly biospecific screening system. *Chemistry & Biodiversity*. 2021
- Gelhaus HC, et al. Efficacy of post exposure administration of doxycycline in a murine model of inhalational melioidosis. *Sci Rep*. 2013;3:1146.
- Xin Y, Chen X, Tang X, et al. Mechanics and actomyosin-dependent survival and chemoresistance of suspended tumor cells in shear flow. *Biophysical Journal*. 2019 May 21;116(10):1803-1814
- Li Y, Qiao K, Zhang X, et al. Targeting myeloid-derived suppressor cells to attenuate vasculogenic mimicry and synergistically enhance the anti-tumor effect of PD-1 inhibitor. *Iscience*. 2021, 24(12): 103392.
- Chen X, Fan Y, Sun J, et al. Nanoparticle-mediated specific elimination of soft cancer stem cells by targeting low cell stiffness. *Acta Biomaterialia*. 2021
- Han L, Jiang Y, Shi M, et al. LIPH contributes to glycolytic phenotype in pancreatic ductal adenocarcinoma by activating LPA/LPAR axis and maintaining ALDOA stability. *Journal of Translational Medicine*. 2023, 21(1): 1-21.
- Hu Z, Lu Y, Cao J, et al. N-acetyltransferase NAT10 controls cell fates via connecting mRNA cytidine acetylation to chromatin signaling. *Science Advances*. 2024, 10(2): eadh9871.

**Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins**

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E\_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481