


## Chrysin

## Chemical Properties

CAS No. :	480-40-0
Formula:	C <sub>15</sub> H <sub>10</sub> O <sub>4</sub>
Molecular Weight:	254.24
Storage:	Powder: -20°C for 3 years   In solvent: -80°C for 1 year Actual storage temperature shall be subject to the COA.



## Biological Description

Description	Chrysin (5,7-Dihydroxyflavone) exhibits many biological activities and pharmacological effects, including antioxidant, anti-inflammatory, anticancer, and antiviral activities.
Targets(IC50)	Estrogen Receptor/ERR,Antioxidant

## Solubility Information

Solubility	Ethanol: < 1 mg/mL (insoluble or slightly soluble), H <sub>2</sub> O: < 1 mg/mL (insoluble or slightly soluble), DMSO: 130 mg/mL (511.33 mM),Sonication is recommended. (< 1 mg/ml refers to the product slightly soluble or insoluble)
In vivo Formulation	10% DMSO+40% PEG300+5% Tween 80+45% Saline: 2 mg/mL (7.87 mM),Sonication is recommended. <i>Please add the solvents sequentially, clarifying the solution as much as possible before adding the next one. Dissolve by heating and/or sonication if necessary. Working solution is recommended to be prepared and used immediately. The formulation provided above is for reference purposes only. In vivo formulations may vary and should be modified based on specific experimental conditions.</i>

## Preparing Stock Solutions

	1mg	5mg	10mg
1 mM	3.9333 mL	19.6665 mL	39.3329 mL
5 mM	0.7867 mL	3.9333 mL	7.8666 mL
10 mM	0.3933 mL	1.9666 mL	3.9333 mL
50 mM	0.0787 mL	0.3933 mL	0.7867 mL

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. Please use it as soon as possible.

Note: The dilution table applies only to solid products. For liquid products, please calculate the stock solution based on the stated concentration and/or density.

### Reference

- Mani R , Natesan V . Chrysin: Sources, beneficial pharmacological activities, and molecular mechanism of action[J]. *Phytochemistry*, 2018, 145:187-196.
- Wang, Ke, Cong Wang, Chuan-Jun Zhu, Ge Li, Yue Li, Yu-Bin Feng, Jing-Jing Ruan et al. 4-Amino-2-Trifluoromethyl-Phenyl Retinate induced leukemia cell differentiation by decreasing eIF6. *Biochemical and Biophysical Research Communications*. 2018, 503(3): 2033-2039
- Liang Y, Xu Z, Wu X, et al. Inhibition of hyperpolarization-activated cyclic nucleotide-gated channels with natural flavonoid quercetin. *Biochemical and Biophysical Research Communications*. 2020
- Wang, Ke, Cong Wang, Chuan-Jun Zhu, Ge Li, Yue Li, Yu-Bin Feng, Jing-Jing Ruan et al. 4-Amino-2-Trifluoromethyl-Phenyl Retinate induced leukemia cell differentiation by decreasing eIF6 [J]. *Biochemical and biophysical research communications*. 2018 Sep 10;503(3):2033-2039.
- Liang Y, Xu Z, Wu X, et al. Inhibition of hyperpolarization-activated cyclic nucleotide-gated channels with natural flavonoid quercetin[J]. *Biochemical and Biophysical Research Communications*. 2020

**Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins**

This product is for Research Use Only· Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E\_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481