

## Cucurbit[8]uril

## Chemical Properties

CAS No. :	259886-51-6
Formula:	C <sub>48</sub> H <sub>48</sub> N <sub>3</sub> O <sub>16</sub>
Molecular Weight:	1329.1
Storage:	Powder: -20°C for 3 years   In solvent: -80°C for 1 year <small>Actual storage temperature shall be subject to the COA.</small>

## Biological Description

Description	Cucurbit[8]uril is a highly effective and safe supramolecular compound that promotes protein heterodimerization. It selectively induces the heterodimerization of methylviologen and naphthalene functionalized proteins, demonstrating its specificity and versatility. Additionally, Cucurbit[8]uril shows exceptional oral activity and low toxicity, making it a promising candidate for various pharmaceutical and biotechnological applications.
Targets(IC50)	Others
In vitro	Cucurbit[8]uril (0~20 μM; 48 hours; CHO-K1 cells) marginally reduces cell viability to 86% [2]. It selectively induces heterodimerization of MV-eYFP with Np-eCFP and high energy transfer between the proteins, observed only with all three supramolecular components forming the ternary complex. Cucurbit[8]uril inhibits unspecific protein assembly induced by methylviologen and enables the formation of selective protein heterodimers with more hydrophobic proteins. It prevents MV-induced unspecific dimerization with hydrophobic protein surfaces[1].
In vivo	Cucurbit[8]uril exhibits minimal toxicity when administered either intravenously or orally in mice, as demonstrated by in vivo studies[2].

## Solubility Information

Solubility	DMSO: < 1 mg/mL (< 1 mg/ml refers to the product slightly soluble or insoluble)
------------	--

### Preparing Stock Solutions

---

	1mg	5mg	10mg
1 mM	0.7524 mL	3.7619 mL	7.5239 mL
5 mM	0.1505 mL	0.7524 mL	1.5048 mL
10 mM	0.0752 mL	0.3762 mL	0.7524 mL
50 mM	0.015 mL	0.0752 mL	0.1505 mL

---

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. Please use it as soon as possible.

Note: The dilution table applies only to solid products. For liquid products, please calculate the stock solution based on the stated concentration and/or density.

### Reference

Uhlenheuer DA, et al. Cucurbit[8]uril induced heterodimerization of methylviologen and naphthalene functionalized proteins. Chem Commun (Camb). 2011;47(24):6798-6800.

Uzunova VD, et al. Toxicity of cucurbit[7]uril and cucurbit[8]uril: an exploratory in vitro and in vivo study. Org Biomol Chem. 2010;8(9):2037-2042.

**Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins**

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E\_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481