

$\alpha$ -Lipoic Acid.

## Chemical Properties


CAS No. : 62-46-4

Formula: C<sub>8</sub>H<sub>14</sub>O<sub>2</sub>S<sub>2</sub>

Molecular Weight: 206.32

Storage: Powder: -20°C for 3 years | In solvent: -80°C for 1 year

Actual storage temperature shall be subject to the COA.



## Biological Description

Description	$\alpha$ -Lipoic Acid (Alphalipoic acid) inhibits NF- $\kappa$ B-dependent HIV-1 LTR activation. $\alpha$ -Lipoic Acid induces endoplasmic reticulum (ER) stress-mediated apoptosis in hepatoma cells. $\alpha$ -Lipoic Acid is an antioxidant, which is an essential cofactor of mitochondrial enzyme complexes.
Targets(IC50)	NF- $\kappa$ B,HIV Protease,Mitochondrial Metabolism,Endogenous Metabolite
In vitro	$\alpha$ -Lipoic Acid (Alpha-Lipoic acid, ALA) is a naturally occurring dithiol compound, plays an essential role in mitochondrial bioenergetics.?It reduces lipid accumulation in the liver by regulating the transcriptional factors SREBP-1, FoxO1, and Nrf2, and their downstream lipogenic targets via the activation of the SIRT1/LKB1/AMPK pathway.? Treatment of cells with $\alpha$ -Lipoic Acid (250, 500 and 1000 $\mu$ M) significantly increases the NAD <sup>+</sup> /NADH ratio in HepG2 cells (P<0.05 or P<0.01).?Treatment with $\alpha$ -Lipoic Acid (50, 125, 250 and 500 $\mu$ M) increases SIRT1 activity in HepG2 cells.? $\alpha$ -Lipoic Acid (50, 125, 250, 500 and 1000 $\mu$ M) increases phosphorylation of AMPK and acetyl-CoA carboxylase (ACC) in HepG2 cells in a dose-dependent fashion
In vivo	Administration of $\alpha$ -Lipoic Acid (100 mg/kg or 200 mg/kg) markedly reduces visceral fat mass in mice.?In addition, $\alpha$ -Lipoic Acid (100 mg/kg or 200 mg/kg) treatment inhibits the appetite and causes a dramatic weight loss (all P<0.05)[1]
Animal Research	C57BL/6J mice, divided into four groups, are fed an high-fat diet (HFD) for 24 weeks to induce nonalcoholic fatty liver disease (NAFLD) followed by daily administration of $\alpha$ -Lipoic Acid.?Then, the effects of $\alpha$ -Lipoic Acid on hepatic lipid accumulation in long-term HFD-fed mice are assessed[1].

## Solubility Information

Solubility	DMSO: 250 mg/mL (1211.71 mM),Sonication is recommended. (< 1 mg/ml refers to the product slightly soluble or insoluble)
------------	--

### Preparing Stock Solutions

---

	1mg	5mg	10mg
1 mM	4.8468 mL	24.2342 mL	48.4684 mL
5 mM	0.9694 mL	4.8468 mL	9.6937 mL
10 mM	0.4847 mL	2.4234 mL	4.8468 mL
50 mM	0.0969 mL	0.4847 mL	0.9694 mL

---

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. Please use it as soon as possible.

Note: The dilution table applies only to solid products. For liquid products, please calculate the stock solution based on the stated concentration and/or density.

### Reference

Xiao L, et al. Activity of the dietary antioxidant ergothioneine in a virus gene-based assay for inhibitors of HIV transcription. *Biofactors*. 2006;27(1-4):157-65.

Pibiri M, et al.  $\alpha$ -Lipoic acid induces Endoplasmic Reticulum stress-mediated apoptosis in hepatoma cells. *Sci Rep*. 2020 Apr 28;10(1):7139.

Yang Y, et al. Alpha-lipoic acid improves high-fat diet-induced hepatic steatosis by modulating the transcription factors SREBP-1, FoxO1 and Nrf2 via the SIRT1/LKB1/AMPK pathway. *J Nutr Biochem*. 2014 Nov;25(11):1207-1217.

**Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins**

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E\_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481