


## Atractylenolide II

## Chemical Properties

CAS No. :	73069-14-4
Formula:	C <sub>15</sub> H <sub>20</sub> O <sub>2</sub>
Molecular Weight:	232.32
Storage:	Keep away from direct sunlight Powder: -20°C for 3 years   In solvent: -80°C for 1 year <small>Actual storage temperature shall be subject to the COA.</small>



## Biological Description

Description	1. Atractylenolide II (2-Atractylenolide) has antimelanoma effect by inhibiting STAT3 signalling. 2. Atractylenolide II has cytotoxic/apoptotic effects may via p38 activation , ERK and Akt inactivation, p53 dependent.
Targets(IC50)	Apoptosis,ERK,Akt,STAT,p38 MAPK,Src

## Solubility Information

Solubility	DMSO: 85 mg/mL (365.87 mM),Sonication is recommended. (< 1 mg/ml refers to the product slightly soluble or insoluble)
In vivo Formulation	10% DMSO+40% PEG300+5% Tween 80+45% Saline: 2 mg/mL (8.61 mM),Sonication is recommended. <i>Please add the solvents sequentially, clarifying the solution as much as possible before adding the next one. Dissolve by heating and/or sonication if necessary. Working solution is recommended to be prepared and used immediately. The formulation provided above is for reference purposes only. In vivo formulations may vary and should be modified based on specific experimental conditions.</i>

## Preparing Stock Solutions

	1mg	5mg	10mg
1 mM	4.3044 mL	21.522 mL	43.0441 mL
5 mM	0.8609 mL	4.3044 mL	8.6088 mL
10 mM	0.4304 mL	2.1522 mL	4.3044 mL
50 mM	0.0861 mL	0.4304 mL	0.8609 mL

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. Please use it as soon as possible.

Note: The dilution table applies only to solid products. For liquid products, please calculate the stock solution based on the stated concentration and/or density.

### Reference

- Fu X Q , Chou G X , Kwan H Y , et al. Inhibition of STAT3 signalling contributes to the antimelanoma action of atractylenolide II[J]. *Experimental Dermatology*, 2014, 23(11):855-857.
- Wang F, Li Z, Chen L, et al. Inhibition of ASCT2 induces hepatic stellate cell senescence with modified proinflammatory secretome through an IL-1 $\alpha$ /NF- $\kappa$ B feedback pathway to inhibit liver fibrosis. *Acta Pharmaceutica Sinica B*. 2022
- Cao L, Xu E, Zheng R, et al. Traditional Chinese medicine Lingguizhugan decoction ameliorate HFD-induced hepatic-lipid deposition in mice by inhibiting STING-mediated inflammation in macrophages. *Chinese Medicine*. 2022, 17(1): 1-16.

**Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins**

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E\_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481