


Ro-3306

Chemical Properties

CAS No. : 872573-93-8
 Formula: C₁₈H₁₃N₃O₂
 Molecular Weight: 351.45
 Storage: Powder: -20°C for 3 years | In solvent: -80°C for 1 year
 Actual storage temperature shall be subject to the COA.



Biological Description

Description	RO-3306 is a selectivity ATP-competitive CDK1 inhibitor (K _i : 20 nM). The selectivity of RO-3306 for CDK1 is >15-fold higher than a diverse panel of human kinases.
Targets(IC50)	Apoptosis,ERK,CDK,PKA,PKC,SGK
In vitro	<p>METHODS: OVCA-429 and OVCAR-3 ovarian cancer cells were treated with Ro 3306 (1-2.5 nM) for 9 days, and cell viability was measured using crystal violet assay.</p> <p>RESULTS: OVCA-429 and OVCAR-3 cells showed low growth rates at high dose concentrations of Ro 3306 for up to 9 days. Treatment with 2.5 μM Ro 3306 reduced the growth rate of OVCA-429 and OVCAR-3 cells by 75.3% and 87.7%, respectively, on day 9. [1]</p> <p>METHODS: Human tumor cell lines HCT116, SW480 and HeLa were treated with Ro 3306 (9 μM) for 20 h. Cell cycle was detected by Flow cytometry.</p> <p>RESULTS: Treatment of proliferating human tumor cells with Ro 3306 for 20 h resulted in complete G2/M cell cycle arrest. [2]</p>
In vivo	In HCT116, SW480, and HeLa cell lines, treatment with RO-3306 for 20 hours resulted in a complete cell cycle arrest at the G2/M phase. RO-3306 (10 μM) effectively inhibits oocyte maturation. The compound demonstrates inhibition of CDK1/cyclin B1 (K _i : 35 nM), CDK1/cyclin A (K _i : 110 nM), CDK2/cyclin E (K _i : 340 nM), and CDK4/cyclin D (K _i >2000 nM) activities. Additionally, RO-3306 significantly impedes the proliferation of HCT116 and SW480 cells. Compared to non-tumorigenic cells (MCF 10A and MCF 12A), RO-3306 exhibits a stronger pro-apoptotic effect on cancerous cells (HCT116 and SW480).
Kinase Assay	CDK assay: The activity of CDK1 cyclin B1, CDK1 cyclin A, CDK2 cyclin E, and CDK4 cyclin D is measured by a homogeneous time-resolved fluorescence assay in a 96-well format. The assay buffer contained 25 mM Hepes, 6.25 mM MgCl ₂ , 0.003% Tween 20, 0.3 mg/mL BSA, 1.5 mM DTT, and ATP as follows: 162 μM (CDK1), 90 μM (CDK2), or 135 μM (CDK4). CDK1 and CDK2 buffer contained 10 mM MgCl ₂ . Test compounds are diluted in assay buffer to 3-fold their final concentration in 20 μL, and the reaction is started by the addition of a 40 μL assay buffer containing the pRB substrate (0.185 μM). The plates are incubated at 37°C for 30 min with constant agitation, and the reaction is terminated by the addition of 15 μL of 1.6 μM anti-phospho pRB antibody (Ser-780) in 25 mM Hepes, 24 mM EDTA, and 0.2 mg/mL BSA. After an additional 30 min of incubation with shaking, 15 μL of 3 nM Lance-Eu-W1024-labeled anti-rabbit IgG and 60 nM Allophycocyanin-conjugated anti-His-6 antibody in 25 mM Hepes, and 0.5 mg/mL BSA is added and

A DRUG SCREENING EXPERT

Kinase Assay	incubated for 1 h. The plates are read in the Victor-V multi- label reader at excitation 340 nm and emission 615 nm and 665 nm. The IC50 values are calculated from the readings at 665 nm and normalized for Europium readings at 615 nm. Ki values are calculated according to the equation: $K_i = IC_{50} / (1 + S/K_m)$, where S is the ATP concentration in the assay and K_m is the Michaelis-Menten constant for ATP. The inhibitory activity against the panel of kinases is determined by the IMAP assay
Cell Research	Log phases cells (25,000) are seeded in 96-well plates and incubated in a 37°C incubator with CO ₂ . After 24 h, different concentrations of RO-3306 are administered to determine the drug concentrations required to achieve a 50% growth inhibition (IC ₅₀). MTT (20 µL, 5 mg/mL stock solution in saline) is added to each well and the cells are incubated for 4 h. Supernatants are removed and formazan crystals from viable cells are solubilized with 200 µL anhydrous DMSO. The absorbance is detected with a 550 model microplate reader at the 565 nm wavelength.(Only for Reference)

Solubility Information

Solubility	DMSO: 22.73 mg/mL (64.67 mM),Sonication is recommended. (< 1 mg/ml refers to the product slightly soluble or insoluble)
In vivo Formulation	10% DMSO+90% (20% SBE-β-CD in Saline): < 2.27 mg/mL (6.46 mM),Lower concentrations may be soluble, but exact solubility limit is unknown. 10% DMSO+90% Saline: < 2.27 mg/mL (6.46 mM),Lower concentrations may be soluble, but exact solubility limit is unknown. 10% DMSO+90% Corn oil: 2.27 mg/mL (6.46 mM),Solution. 10% DMSO+40% PEG300+5% Tween 80+45% Saline: 1.25 mg/mL (3.56 mM),Solution. <i>Please add the solvents sequentially, clarifying the solution as much as possible before adding the next one. Dissolve by heating and/or sonication if necessary. Working solution is recommended to be prepared and used immediately. The formulation provided above is for reference purposes only. In vivo formulations may vary and should be modified based on specific experimental conditions.</i>

Preparing Stock Solutions

	1mg	5mg	10mg
1 mM	2.8454 mL	14.2268 mL	28.4535 mL
5 mM	0.5691 mL	2.8454 mL	5.6907 mL
10 mM	0.2845 mL	1.4227 mL	2.8454 mL
50 mM	0.0569 mL	0.2845 mL	0.5691 mL

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. Please use it as soon as possible.

Note: The dilution table applies only to solid products. For liquid products, please calculate the stock solution based on the stated concentration and/or density.

Reference

- Yang W, et al. Accumulation of cytoplasmic Cdk1 is associated with cancer growth and survival rate in epithelial ovarian cancer. *Oncotarget*. 2016 Aug 2;7(31):49481-49497.
- Zhang Y, Hong X, Hua S, et al. Reconstitution and mechanistic dissection of the human microtubule branching machinery. *Journal of Cell Biology*. 2022, 221(7): e202109053
- Chen C, Zhang X, Wang Y, et al. Translational and Post-translational Control of Human Naïve versus Primed Pluripotency. *iScience*. 2021: 103645.
- Vassilev LT, et al. Selective small-molecule inhibitor reveals critical mitotic functions of human CDK1. *Proc Natl Acad Sci U S A*. 2006 Jul 11;103(28):10660-5.
- Xia Q, et al. *Int J Oncol*. 2014, 44(3), 735-744.
- Ou J, Li H, Qiu P, et al. CDK9 modulates circadian clock by attenuating REV-ERB α activity. *Biochemical and Biophysical Research Communications*. 2019 Jun 11;513(4):967-973
- Ou J, Li H, Qiu P, et al. CDK9 modulates circadian clock by attenuating REV-ERB α activity[J]. *Biochemical and biophysical research communications*. 2019 Jun 11;513(4):967-973.
- Guo S, Lei X, Chang Y, et al. SARS-CoV-2 hijacks cellular kinase CDK2 to promote viral RNA synthesis. *Signal Transduction and Targeted Therapy*. 2022, 7(1): 1-12.
- Jiang L, Yu Y, Li Z, et al. BMS-265246, a Cyclin-Dependent Kinase Inhibitor, Inhibits the Infection of Herpes Simplex Virus Type 1. *Viruses*. 2023, 15(8): 1642.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel: 781-999-4286 E_mail: info@targetmol.com Address: 34 Washington Street, Wellesley Hills, MA 02481