


Lapatinib Ditosylate

Chemical Properties

CAS No. :	388082-77-7
Formula:	C ₂₉ H ₂₆ ClFN ₄ O ₄ S·2C ₇ H ₈ O ₃ S
Molecular Weight:	925.46
Storage:	Powder: -20°C for 3 years In solvent: -80°C for 1 year Actual storage temperature shall be subject to the COA.



Biological Description

Description	Lapatinib Ditosylate (Tykerb ditosylate) is an effective EGFR and ErbB2 inhibitor (IC ₅₀ : 10.8/9.2 nM for EGFR/ErbB2).
Targets(IC ₅₀)	EGFR, Ferroptosis, Autophagy
In vitro	Lapatinib (1 μM) induces apoptosis in NCI-N87 and OD19 cells [2]. Lapatinib inhibits the growth of EGFR-overexpressing A431 skin cancer (IC ₅₀ : 0.14 μM) and ErbB2-overexpressing SK-BR-3 breast cancer cells (IC ₅₀ : 0.124 μM). It also inhibits the growth of ErbB2-amplified OD19 esophageal (IC ₅₀ : 0.09 μM) and NCI-N87 gastric cancer cells (IC ₅₀ : 0.01 μM) as well as several types of gastric cancer cells in which ErbB2 is not amplified (IC ₅₀ s: 0.35-8.58 μM) [3].
In vivo	Lapatinib (30 and 100 mg/kg, p.o., b.i.d) dose-responsive inhibited the growth of BT474 and HN5 human tumor xenografts. Complete inhibition of tumor growth is seen at the 100 mg/kg dose. At this dose, there is <10% weight loss in treated animals over the course of the 21-day treatment. Lapatinib treatment inhibits tumor xenograft growth of the HN5 and BT474 cells in a dose-responsive manner at 30 and 100 mg/kg orally, twice daily, with complete inhibition of tumor growth at the higher dose [1]. Lapatinib (100 mg/kg/day, p.o.) induces severe oxidative damage in the cardiac tissue of rat [4].
Kinase Assay	The IC ₅₀ values for inhibition of enzyme activity are generated by measuring the inhibition of phosphorylation of a peptide substrate. The intracellular kinase domains of EGFR and ErbB2 are purified from a baculovirus expression system. EGFR and ErbB2 reactions are performed in 96-well polystyrene round-bottomed plates in a final volume of 45 μL. Reaction mixtures contain 50 mM 4-morpholinepropanesulfonic acid (pH 7.5), 2 mM MnCl ₂ , 10 μM ATP, 1 μCi of [³³ P] ATP/reaction, 50 μM Peptide A [Biotin-(amino hexanoic acid)-EEEEYFELVAKKK-CONH ₂], 1 mM dithiothreitol, and 1 μL of DMSO containing serial dilutions of Lapatinib beginning at 10 μM. The reaction is initiated by adding the indicated purified type-1 receptor intracellular domain. The amount of enzyme added is 1 pmol/reaction (20 nM). Reactions are terminated after 10 minutes at 23°C by adding 45 μL of 0.5% phosphoric acid in water. The terminated reaction mix (75 μL) is transferred to phosphocellulose filter plates. The plates are filtered and washed three times with 200 μL of 0.5% phosphoric acid. Scintillation cocktail (50 μL) is added to each well, and the assay is quantified by counting in a Packard Topcount. IC ₅₀ values are generated from 10-point dose-response curves [1].

Cell Research	Cells are plated in 96-well plates, in the media, at the following densities: HFF and HN5, 1000 cells/well and BT474, 5000 cells/well. After 24 h, the cells are exposed to vehicle (0.3% DMSO) or Lapatinib (1 nM, 10 nM, 100 nM, 1 μM, 10 μM, and 100 μM). Lapatinib is removed from the cells after 72 h and is replaced by either DMEM containing 10% FBS and 50 μg/mL Gentamicin (HFF and HN5) or RPMI containing 10% FBS and 50 μg/mL Gentamicin (BT474). Methylene blue staining is performed at the time points over a total period of 16 days. Relative cell number is estimated using methylene blue staining. The absorbance at 620 nm is read in a Spectra microplate reader. Cell death and cell cycle analysis are assessed by propidium iodide staining and antibody detection of incorporated BrdUrd and staining with propidium iodide [1].
Animal Research	CD-1 nude female mice are used for HN5 human tumor xenografts, which are initiated by injection of a cell suspension in PBS: Matrigel (1:1). C.B-17 SCID female mice are used for BT474 human tumor xenografts, which are initiated by implantation of tumor fragments (20-100 mg) from established tumors. Tumor cells and fragments are implanted by s.c. injection in the right flank. The s.c. tumors are measured with calipers, and mice are weighed twice weekly. Tumor weight is estimated from tumor volume using this formula: length×width ² /2=tumor volume (mm ³). Treatment begins when tumors are palpable, 3-5 mm in diameter. Lapatinib (30 and 100 mg/kg) is administered p.o. twice daily for 21 days in a vehicle of sulfo-butyl-ether-β-cyclodextrin 10% aqueous solution (CD10) [1].

Solubility Information

Solubility	Ethanol: < 1 mg/mL (insoluble or slightly soluble), DMSO: 250 mg/mL (270.14 mM),Sonication is recommended. H2O: < 1 mg/mL (insoluble or slightly soluble), (< 1 mg/ml refers to the product slightly soluble or insoluble)
In vivo Formulation	10% DMSO+40% PEG300+5% Tween 80+45% Saline: 3.3 mg/mL (3.57 mM),Sonication is recommended. <i>Please add the solvents sequentially, clarifying the solution as much as possible before adding the next one. Dissolve by heating and/or sonication if necessary. Working solution is recommended to be prepared and used immediately. The formulation provided above is for reference purposes only. In vivo formulations may vary and should be modified based on specific experimental conditions.</i>

Preparing Stock Solutions

	1mg	5mg	10mg
1 mM	1.0805 mL	5.4027 mL	10.8054 mL
5 mM	0.2161 mL	1.0805 mL	2.1611 mL
10 mM	0.1081 mL	0.5403 mL	1.0805 mL
50 mM	0.0216 mL	0.1081 mL	0.2161 mL

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. Please use it as soon as possible.

Note: The dilution table applies only to solid products. For liquid products, please calculate the stock solution based on the stated concentration and/or density.

Reference

Rusnak DW, et al. The effects of the novel, reversible epidermal growth factor receptor/ErbB-2 tyrosine kinase inhibitor, GW2016, on the growth of human normal and tumor-derived cell lines in vitro and in vivo. *Mol Cancer Ther.* 2001 Dec;1(2):85-94

Chefrour M et al. Positive interaction between lapatinib and capecitabine in human breast cancer models: study of molecular determinants. *Fundam Clin Pharmacol.* 2012 Aug;26(4):530-7.

Wainberg ZA, et al. Lapatinib, a dual EGFR and HER2 kinase inhibitor, selectively inhibits HER2-amplified human gastric cancer cells and is synergistic with trastuzumab in vitro and in vivo. *Clin Cancer Res.* 2010 Mar 1;16(5):1509-19.

Eryilmaz U, et al. S100A1 as a Potential Diagnostic Biomarker for Assessing Cardiotoxicity and Implications for the Chemotherapy of Certain Cancers. *PLoS One.* 2015 Dec 18;10(12):e0145418.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only· Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481