

Sotuletinib


Chemical Properties

CAS No. : 953769-46-5

Formula: C₂₀H₂₂N₄O₃S

Molecular Weight: 398.48

Storage: Keep away from direct sunlight, Keep away from moisture, Store at low temperature
 Powder: -20°C for 3 years | In solvent: -80°C for 1 year
Actual storage temperature shall be subject to the COA.



Biological Description

Description	Sotuletinib (BLZ945) is an orally active, effective and specific CSF-1R inhibitor (IC ₅₀ : 1 nM), >1000-fold selective against its closest receptor tyrosine kinase homologs.
Targets(IC ₅₀)	c-Fms, CSF-1R
In vitro	In bone marrow-derived macrophages (BMDMs), Sotuletinib specifically inhibits CSF-1-dependent proliferation with EC ₅₀ of 67 nM, and decreases CSF-1R phosphorylation. Sotuletinib blocks the reciprocal effects between macrophages and glioma cells on the survival, proliferation and/ or polarization of each other to promote tumorigenesis. [1]
In vivo	In glioma-bearing mice, Sotuletinib blocks tumor progression and significantly improves survival via CSF-1R inhibition. Sotuletinib also inhibits orthotopic tumor growth of patient-derived proneural tumor spheres and cell lines in vivo. [1] Sotuletinib (200 mg/kg, p.o.) decreases the growth of malignant cells in both mouse mammary tumor virus-driven polyomavirus middle T antigen (MMTV-PyMT) model of mammary carcinogenesis and keratin 14-expressing human papillomavirus type 16 (K14-HPV-16) transgenic model of cervical carcinogenesis. [2]
Kinase Assay	Inhibition of biochemical TrkA, TrkB and TrkC: TrkA and TrkC biochemical assays are carried out by HTRF method. The reaction mixtures contains 1 μM peptide substrate, 1 μM ATP, and either 1.8 nM TrkA or 34 nM TrkC in the reaction buffer (50 mM HEPES pH 7.1, 10 mM MgCl ₂ , 2 mM MnCl ₂ , 0.01% BSA, 2.5 mM DTT and 0.1 mM Na ₃ VO ₄) at a final volume of 10 μL. All reactions are carried out at room temperature in white ProxiPlate ² 384-well Plus plates and are quenched with 5 μL of 0.2 mM EDTA at 60 min. Five μL of the detection reagents (2.5 ng PT66K and 0.05 μg SAXL per well) are added, the plates are incubated at room temperature for 1 h and then read in EnVision reader. Compounds are diluted into assay mixture (final DMSO 0.5%), and IC ₅₀ values are determined by 12-point (from 50 to 0.000282 μM) inhibition curves in duplicate under the assay conditions. TrkB biochemical assay is carried out by caliper microfluidic method. The reaction mixtures contained 1 μM peptide substrate, 10 μM ATP, and 2 nM TrkB in a reaction buffer containing 100 mM HEPES, pH 7.5, 5 mM MgCl ₂ , 0.01% Triton X-100, 0.1% BSA, 1 mM DTT, 10 μM Na ₃ VO ₄ , and 10 μM Beta-Glycerophosphate. The reactions are carried out at room temperature for 3 hrs, and the products are determined by Caliper EZ-reader. Compounds are diluted into assay mixture (final DMSO

A DRUG SCREENING EXPERT

Kinase Assay	1%), and IC50 values are determined by 12-point (from 50 to 0.000282 μ M) inhibition curves in duplicate under the assay conditions.
Cell Research	Cell growth rate is determined using the MTT cell proliferation kit. Briefly, cells are plated in triplicate in 96-well plates: 1,000 cells per well for glioma cell lines, 5 x 1,000 cells per well for BMDM and CRL-2467, and 2.5 x 1,000 cells per well for HUVEC and HBMEC cell lines. For all experiments, media is changed every 48 h. Cells are grown in the presence or absence of 6.7–6,700 nM of BLZ945, or 8 μ g/mL of CSF-1R neutralizing antibody. BMDM and CRL-2467 cells were supplemented with 10 ng/mL and 30 ng/mL recombinant mouse CSF-1, respectively. Reduction of the MTT substrate is detected by colorimetric analysis using a plate reader as per the manufacturer's protocol, and measured at 595 nm and 750 nm on a spectraMax 340pc plate reader.(Only for Reference)

Solubility Information

Solubility	DMSO: 257 mg/mL (644.95 mM),Sonication is recommended. Ethanol: 3 mg/mL (7.53 mM),Heating is recommended. H2O: < 1 mg/mL (insoluble or slightly soluble), (< 1 mg/ml refers to the product slightly soluble or insoluble)
In vivo Formulation	10% DMSO+40% PEG300+5% Tween 80+45% Saline: 5 mg/mL (12.55 mM),Sonication is recommended. <i>Please add the solvents sequentially, clarifying the solution as much as possible before adding the next one. Dissolve by heating and/or sonication if necessary. Working solution is recommended to be prepared and used immediately. The formulation provided above is for reference purposes only. In vivo formulations may vary and should be modified based on specific experimental conditions.</i>

Preparing Stock Solutions

	1mg	5mg	10mg
1 mM	2.5095 mL	12.5477 mL	25.0954 mL
5 mM	0.5019 mL	2.5095 mL	5.0191 mL
10 mM	0.251 mL	1.2548 mL	2.5095 mL
50 mM	0.0502 mL	0.251 mL	0.5019 mL

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. Please use it as soon as possible.

Note: The dilution table applies only to solid products. For liquid products, please calculate the stock solution based on the stated concentration and/or density.

Reference

Pyonteck SM, et al. Nat Med. 2013, 19(10), 1264-1272.

Fan K, Li Y, Wang H, et al. Stress-Induced Metabolic Disorder in Peripheral CD4+ T Cells Leads to Anxiety-like Behavior. Cell. 2019, 179(4): 864-879. e19.

Strachan DC, et al. Oncoimmunology. 2013, 2(12), e26968.

Huang C, Ou R, Chen X, et al. Tumor cell-derived SPON2 promotes M2-polarized tumor-associated macrophage infiltration and cancer progression by activating PYK2 in CRC. Journal of Experimental & Clinical Cancer Research. 2021, 40(1): 1-17.

Fan K, Li Y, Wang H, et al. Stress-Induced Metabolic Disorder in Peripheral CD4+ T Cells Leads to Anxiety-like Behavior[J]. Cell. 2019, 179(4): 864-879. e19.

Haydar D, Ibañez-Vega J, Crawford J C, et al. CAR T-cell design-dependent remodeling of the brain tumor-immune microenvironment modulates tumor-associated macrophages and anti-glioma activity. Cancer Research Communications. 2023

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel: 781-999-4286 E_mail: info@targetmol.com Address: 34 Washington Street, Wellesley Hills, MA 02481