


Oridonin

Chemical Properties

CAS No. :	28957-04-2
Formula:	C ₂₀ H ₂₈ O ₆
Molecular Weight:	364.44
Storage:	Powder: -20°C for 3 years In solvent: -80°C for 1 year <i>Actual storage temperature shall be subject to the COA.</i>



Biological Description

Description	Oridonin (NSC-250682) is a diterpenoid isolated from <i>Rabdosia rubescens</i> with anti-tumor, anti-bacterial and anti-inflammatory effects. It acts as an inhibitor of AKT, with IC50s of 8.4 and 8.9 μ M for AKT1 and AKT2.
Targets(IC50)	Akt,Antibacterial
In vivo	LD50: Mice 35-40 mg/kg. [4]

Solubility Information

Solubility	H ₂ O: < 1 mg/mL (insoluble or slightly soluble), Ethanol: 28 mg/mL (76.83 mM),Sonication is recommended. DMSO: 65 mg/mL (178.36 mM),Sonication is recommended. (< 1 mg/ml refers to the product slightly soluble or insoluble)
In vivo Formulation	10% DMSO+40% PEG300+5% Tween 80+45% Saline: 2 mg/mL (5.49 mM),Sonication is recommended. <i>Please add the solvents sequentially, clarifying the solution as much as possible before adding the next one. Dissolve by heating and/or sonication if necessary. Working solution is recommended to be prepared and used immediately. The formulation provided above is for reference purposes only. In vivo formulations may vary and should be modified based on specific experimental conditions.</i>

Preparing Stock Solutions

	1mg	5mg	10mg
1 mM	2.7439 mL	13.7197 mL	27.4394 mL
5 mM	0.5488 mL	2.7439 mL	5.4879 mL
10 mM	0.2744 mL	1.372 mL	2.7439 mL
50 mM	0.0549 mL	0.2744 mL	0.5488 mL

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. Please use it as soon as possible.

Note: The dilution table applies only to solid products. For liquid products, please calculate the stock solution based on the stated concentration and/or density.

Reference

Ikezoe T, et al. *Int J Oncol*, 2003, 23(4), 1187-1193.

Li L, Song J, Zhang M, et al. Oridonin ameliorates caspase-9-mediated brain neuronal apoptosis in mouse with ischemic stroke by inhibiting RIPK3-mediated mitophagy. *Acta Pharmacologica Sinica*. 2022: 1-15.

An Y, Zhu J, Wang X, et al. Oridonin Delays Aging Through the AKT Signaling Pathway. *Frontiers in Pharmacology*. 2022: 1823

Ikezoe T, et al. *Mol Cancer Ther*, 2005, 4(4), 578-586.

Li L, Cheng S Q, Guo W, et al. Oridonin prevents oxidative stress-induced endothelial injury via promoting Nrf-2 pathway in ischaemic stroke. *Journal of Cellular and Molecular Medicine*. 2021

Cui Q, et al. *J Pharmacol Sci*, 2007, 105(4), 317-325.

Yan H, Zhang R, Yan G, et al. Validation of baicalein and oridonin as nonspecific SARS-CoV-2 main protease inhibitors. *Phytotherapy Research*. 2023

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel: 781-999-4286 E_mail: info@targetmol.com Address: 34 Washington Street, Wellesley Hills, MA 02481