

CD28 Protein, Cynomolgus, Recombinant (hFc)

General Information

Synonyms:	CD28 molecule
Protein Construction:	A DNA sequence encoding the cynomolgus CD28 [(Identical to the rhesus CD28 (NP_001036106.2)) (Met1-Pro152)] was expressed with the Fc region of human IgG1 at the C-terminus. Predicted N terminal: Asn 19
Species:	Cynomolgus
Expression Host:	HEK293 Cells
Accession:	Q0PDN3
Molecular Weight:	41.9 kDa (predicted)

QC Testing

Biological Activity:	<ol style="list-style-type: none">1. Immobilized Recombinant Human CD86 / B7-2 Protein (His Tag) at 2 µg/mL (100 µL/well) can bind Recombinant Cynomolgus CD28 / TP44 Protein (Fc Tag, ECD), the EC50 is 100-420 ng/mL.2. Immobilized Recombinant Human CD80 / B7-1 Protein (His Tag) at 2 µg/mL (100 µL/well) can bind Recombinant Cynomolgus CD28 / TP44 Protein (Fc Tag, ECD), the EC50 is 600-1400 ng/mL.
Purity:	> 90 % as determined by SDS-PAGE.
Endotoxin:	< 1.0 EU/µg of the protein as determined by the LAL method.
Formulation:	Lyophilized from a solution filtered through a 0.22 µm filter, containing PBS, pH 7.4. Typically, a mixture containing 5% to 8% trehalose, mannitol, and 0.01% Tween 80 is incorporated as a protective agent before lyophilization.

Preparation and Storage

Reconstitution:

A Certificate of Analysis (CoA) containing reconstitution instructions is included with the products. Please refer to the CoA for detailed information.

Stability & Storage:

It is recommended to store recombinant proteins at -20°C to -80°C for future use. Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Actual storage temperature shall be subject to the COA.

Shipping:

In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

Protein Background

CD28 (Cluster of Differentiation 28) is a disulphide-bonded glycoprotein belonging to the immunoglobulin (Ig) superfamily, and structurally consists of a single Ig V-like extracellular domain, a transmembrane domain and an intracellular domain. Mouse CD28 is constitutively expressed on the surface of all murine T cells and on developing thymocytes as disulfide-linked homodimers or as monomers. CD28 can binds the B7-1 and B7-2 ligand, and together perform important functions in the T and B cell response pathways. B7/CD28 family members, which can augment or antagonize T-cell receptor signaling, in the regulation of central and peripheral T-cell tolerance. CD28 is thus involved in T-cell activation, the induction of cell proliferation and cytokine production and promotion of T-cell survival. Cancer Immunotherapy Co-stimulatory Immune Checkpoint Targets Immune Checkpoint Immune Checkpoint Detection: Antibodies Immune Checkpoint Detection: ELISA Antibodies Immune Checkpoint Detection: FCM Antibodies Immune Checkpoint Detection: IHC Antibodies Immune Checkpoint Detection: WB Antibodies Immune Checkpoint Targets Immunotherapy Targeted Therapy

Reference

- Keir ME, et al. (2005) The B7/CD28 costimulatory family in autoimmunity. *Immunol Rev.* 204: 128-43.
- Sansom DM, et al. (2006) The role of CD28 and cytotoxic T-lymphocyte antigen-4 (CTLA-4) in regulatory T-cell biology. *Immunol Rev.* 212: 131-48.
- Bjrgo E, et al. (2010) Novel mechanism of signaling by CD28. *Immunol Lett.* 129(1): 1-6.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel: 781-999-4286 E_mail: info@targetmol.com Address: 34 Washington Street, Wellesley Hills, MA 02481