

Interferon alpha 2/IFNA2 Protein, Rhesus, Recombinant

General Information

Synonyms:	interferon, alpha 2;interferon, α 2;Interferon α 2/IFNA2
Protein Construction:	A DNA sequence encoding the rhesus IFNA2 (NP_001129266.1) (Cys24-Glu188) was expressed. Predicted N terminal: Cys 24
Species:	Rhesus
Expression Host:	P. pastoris (Yeast)
Accession:	B6CK11
Molecular Weight:	19.5 kDa (predicted)

QC Testing

Biological Activity:	Measured in antiviral assay using WISH cells infected with vesicular stomatitis virus. The ED50 for this effect is typically 1.8-7.4 pg/mL.
Purity:	> 90 % as determined by SDS-PAGE
Endotoxin:	Please contact us for more information.
Formulation:	Lyophilized from a solution filtered through a 0.22 μ m filter, containing PBS, pH 7.4. Typically, a mixture containing 5% to 8% trehalose, mannitol, and 0.01% Tween 80 is incorporated as a protective agent before lyophilization.

Preparation and Storage

Reconstitution:
A Certificate of Analysis (CoA) containing reconstitution instructions is included with the products. Please refer to the CoA for detailed information.

Stability & Storage:

It is recommended to store recombinant proteins at -20°C to -80°C for future use. Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Actual storage temperature shall be subject to the COA.

Shipping:

In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

Protein Background

IFNA2 (Interferon Alpha 2) is a Protein Coding gene. This gene is a member of the alpha interferon gene cluster on chromosome 9. The encoded protein is a cytokine produced in response to viral infection. Type I Interferons (IFNs) are well-known cytokines that exert antiviral activity, antitumor activity, and immunomodulatory effects. Interferon tau (IFNT), a type I IFN similar to alpha IFNs (IFNA), is the pregnancy recognition signal produced by the ruminant conceptus. Among the IFN- α genes, a total of 28 different sequence variants have been described. The three

principal subtypes of IFN α -2 are designated α -2a, α -2b, and α -2c. IFN α -2b is being the predominant allele while IFN α -2a is less predominant and IFN α -2c only a minor allelic variant.

Reference

Wang, et al. (2004) Fever of recombinant human interferon-alpha is mediated by opioid domain interaction with opioid receptor inducing prostaglandin E2. *J Neuroimmunol.* 156(1-2): 107-12.

Groopman JE, et al. (1984) Recombinant alpha-2 interferon therapy for Kaposi's sarcoma associated with the acquired immunodeficiency syndrome. *Ann Intern Med.* 100(5): 671-6.

Krueger JM, et al. (1987) Interferon alpha-2 enhances slow-wave sleep in rabbits. *Int J Immunopharmacol.* 9(1): 23-30.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel: 781-999-4286 E_mail: info@targetmol.com Address: 34 Washington Street, Wellesley Hills, MA 02481