

Influenza A H1N9 (A/mallard/Ohio/265/1987) Hemagglutinin/HA Protein (His)

General Information

Synonyms:	Harvey rat sarcoma viral oncogene homolog
Protein Construction:	A DNA sequence encoding the Influenza A virus(A/mallard/Ohio/265/1987 (H1N9)) hemagglutinin (ABK40634.1)(Met1-Gln529), termed as HA, was expressed with a polyhistidine tag at the C-terminus. Predicted N terminal: Asp 17
Species:	H1N9
Expression Host:	Baculovirus Insect Cells
Accession:	A0M895
Molecular Weight:	58.6 kDa (predicted)

QC Testing

Biological Activity:	Measured by its ability to agglutinate guinea pig red blood cells. HA titer is 0.2-1 μ g/mL for 1% GRBC.
Purity:	> 90 % as determined by SDS-PAGE.
Endotoxin:	< 1.0 EU/ μ g of the protein as determined by the LAL method.
Formulation:	Lyophilized from a solution filtered through a 0.22 μ m filter, containing 20 mM Tris, 500 mM NaCl, 10% glycerol, pH 7.5. Typically, a mixture containing 5% to 8% trehalose, mannitol, and 0.01% Tween 80 is incorporated as a protective agent before lyophilization.

Preparation and Storage

Reconstitution:
A Certificate of Analysis (CoA) containing reconstitution instructions is included with the products. Please refer to the CoA for detailed information.

Stability & Storage:

It is recommended to store recombinant proteins at -20°C to -80°C for future use. Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Actual storage temperature shall be subject to the COA.

Shipping:

In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

Protein Background

The influenza viral Hemagglutinin (HA) protein is a homotrimer with a receptor binding pocket on the globular head of each monomer. HA has at least 18 different antigens. These subtypes are named H1 through H18. HA has two functions. Firstly, it allows the recognition of target vertebrate cells, accomplished through the binding to these cells' sialic acid-containing receptors. Secondly, once bound it facilitates the entry of the viral genome into

the target cells by causing the fusion of the host endosomal membrane with the viral membrane. The influenza virus Hemagglutinin (HA) protein is translated in cells as a single protein, HA, or hemagglutinin precursor protein. For viral activation, hemagglutinin precursor protein (HA) must be cleaved by a trypsin-like serine endoprotease at a specific site, normally coded for by a single basic amino acid (usually arginine) between the HA1 and HA2 domains of the protein. After cleavage, the two disulfide-bonded protein domains produce the mature form of the protein subunits as a prerequisite for the conformational change necessary for fusion and hence viral infectivity.

Reference

White JM, Hoffman LR, Arevalo JH, et al. Attachment and entry of influenza virus into host cells. Pivotal roles of hemagglutinin. In Chiu W, Burnett RM, Garcea RL. Structural Biology of Viruses. 1997 Suzuki Y. Sialobiology of influenza: molecular mechanism of host range variation of influenza viruses. Biol. Pharm. Bull. 2005.

Senne DA, Panigrahy B, Kawaoka Y, et al. Survey of the hemagglutinin (HA) cleavage site sequence of H5 and H7 avian influenza viruses: amino acid sequence at the HA cleavage site as a marker of pathogenicity potential. Avian Dis. 1996 Donald J. Benton, Influenza hemagglutinin membrane anchor, PNAS, 2018

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel: 781-999-4286 E_mail: info@targetmol.com Address: 34 Washington Street, Wellesley Hills, MA 02481