

## VISTA Protein, Rhesus, Recombinant (His)

### General Information

Synonyms:	V-set immunoregulatory receptor
Protein Construction:	A DNA sequence encoding the rhesus C10orf54 (NP_001177807.1) (Met1-Ala194) was expressed with a polyhistidine tag at the C-terminus. Predicted N terminal: Phe 33
Species:	Rhesus
Expression Host:	HEK293 Cells
Accession:	F7F6S6
Molecular Weight:	19.6 kDa (predicted)

### QC Testing

Biological Activity:	Activity testing is in progress. It is theoretically active, but we cannot guarantee it. If you require protein activity, we recommend choosing the eukaryotic expression version first.
Purity:	> 95 % as determined by SDS-PAGE.
Endotoxin:	< 1.0 EU/μg of the protein as determined by the LAL method.
Formulation:	Lyophilized from a solution filtered through a 0.22 μm filter, containing PBS, pH 7.4. Typically, a mixture containing 5% to 8% trehalose, mannitol, and 0.01% Tween 80 is incorporated as a protective agent before lyophilization.

### Preparation and Storage

Reconstitution:	A Certificate of Analysis (CoA) containing reconstitution instructions is included with the products. Please refer to the CoA for detailed information.
Stability & Storage:	It is recommended to store recombinant proteins at -20°C to -80°C for future use. Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots. <small>Actual storage temperature shall be subject to the COA.</small>
Shipping:	In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

### Protein Background

VSIR (V-Set Immunoregulatory Receptor, also known as VISTA) is a Protein Coding gene. VISTA is an immunoregulatory receptor that inhibits the T-cell response. It may promote differentiation of embryonic stem cells, by inhibiting BMP4 signaling. VSIR, or V-set immunoregulatory receptor, could be involved in the pathogenesis of chronic rhinosinusitis with nasal polyps. V-domain Immunoglobulin Suppressor of T cell Activation (VISTA) is an inhibitory immune-checkpoint molecule that suppresses CD4+ and CD8+ T cell activation

## A DRUG SCREENING EXPERT

---

when expressed on antigen-presenting cells. VSIR is broadly expressed in the spleen, bone marrow, and other tissues. Diseases associated with VSIR include Ichthyosis, Congenital, Autosomal Recessive 6, and Monckeberg Arteriosclerosis. An important paralog of this gene is VSIG8.

**Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins**

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E\_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481