

Beta Amylase/AMY2B Protein, Human, Recombinant (hFc)

General Information

Synonyms:	β Amylase/AMY2B;AMY2;HXA;amylase, α 2B (pancreatic);AMY3;amylase, alpha 2B (pancreatic)
Protein Construction:	A DNA sequence encoding the human AMY2B (NP_066188.1) (Met1-Leu511) was expressed with the Fc region of human IgG1 at the C-terminus. Predicted N terminal: Gln 16
Species:	Human
Expression Host:	HEK293 Cells
Accession:	P19961-1
Molecular Weight:	82.4 kDa (predicted)

QC Testing

Biological Activity:	Activity testing is in progress. It is theoretically active, but we cannot guarantee it. If you require protein activity, we recommend choosing the eukaryotic expression version first.
Purity:	> 95 % as determined by SDS-PAGE.
Endotoxin:	< 1.0 EU/ μ g of the protein as determined by the LAL method.
Formulation:	Lyophilized from a solution filtered through a 0.22 μ m filter, containing PBS, pH 7.4. Typically, a mixture containing 5% to 8% trehalose, mannitol, and 0.01% Tween 80 is incorporated as a protective agent before lyophilization.

Preparation and Storage

Reconstitution:

A Certificate of Analysis (CoA) containing reconstitution instructions is included with the products. Please refer to the CoA for detailed information.

Stability & Storage:

It is recommended to store recombinant proteins at -20°C to -80°C for future use. Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Actual storage temperature shall be subject to the COA.

Shipping:

In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

Protein Background

Amylases are secreted proteins that hydrolyze 1,4-alpha-glucoside bonds in oligosaccharides and polysaccharides, and thus catalyze the first step in digestion of dietary starch and glycogen. Alpha-amylase is the major form of amylase found in humans and other mammals. Alpha-amylase hydrolyses alpha bonds of large, alpha-linked polysaccharides, such as starch and glycogen, yielding glucose and maltose. Amylases is widely

expressed and is most prominent in pancreatic juice and saliva, each of which has its own isoform of human α -amylase. They behave differently on isoelectric focusing, and can also be separated in testing by using specific monoclonal antibodies.

Reference

Abe A, et al. (2005) Complexes of *Thermoactinomyces vulgaris* R-47 Alpha-amylase / AMY2A 1 and pullulan model oligosaccharides provide new insight into the mechanism for recognizing substrates with alpha-(1,6) glycosidic linkages. *FEBS J.* 272(23):6145-53.

Aghajari, N, et al. (1998) Crystal structures of the psychrophilic Alpha-amylase / AMY2A from *Alteromonas haloplanctis* in its native form and complexed with an inhibitor. *Protein Sci.* 7(3): 564-72.

Ramasubbu, N, et al. (1996) Structure of Human Salivary α -Amylase at 1.6 Resolution: Implications for its Role in the Oral Cavity. *Acta Crystallographica Section D Biological Crystallography.* 52(3):435-46.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481