

VMO1 Protein, Human, Recombinant (mFc)

General Information

Synonyms:	ERGA6350;PRO21055;vitelline membrane outer layer 1 homolog (chicken)
Protein Construction:	A DNA sequence encoding the human VMO1 (NP_872372.1) (Met1-Ser202) was expressed with the Fc region of mouse IgG1 at the C-terminus. Predicted N terminal: Gln 25
Species:	Human
Expression Host:	HEK293 Cells
Accession:	Q7Z5L0-1
Molecular Weight:	45.4 kDa (predicted)

QC Testing

Biological Activity:	Activity testing is in progress. It is theoretically active, but we cannot guarantee it. If you require protein activity, we recommend choosing the eukaryotic expression version first.
Purity:	> 80 % as determined by SDS-PAGE.
Endotoxin:	< 1.0 EU/μg of the protein as determined by the LAL method.
Formulation:	Lyophilized from a solution filtered through a 0.22 μm filter, containing PBS, pH 7.4. Typically, a mixture containing 5% to 8% trehalose, mannitol, and 0.01% Tween 80 is incorporated as a protective agent before lyophilization.

Preparation and Storage

Reconstitution:	A Certificate of Analysis (CoA) containing reconstitution instructions is included with the products. Please refer to the CoA for detailed information.
Stability & Storage:	It is recommended to store recombinant proteins at -20°C to -80°C for future use. Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots. <small>Actual storage temperature shall be subject to the COA.</small>
Shipping:	In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

Protein Background

Vitelline membrane outer layer protein 1 (VMO1), a basic protein present in the outer layer of the vitelline membrane of eggs, plays essential roles in separating the yolk from the egg white and preventing infection from bacteria by forming a barrier of fibrous layers in avian eggs. VMO1 is regulated by estrogen and target microRNAs and it is a potential diagnostic marker of ovarian cancer in laying hens.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481