

Notch 1 Protein, Mouse, Recombinant (hFc)

General Information

Synonyms:	lin-12;Tan1;Mis6;9930111A19Rik;notch 1;N1;Notch1
Protein Construction:	A DNA sequence encoding the mouse Notch1 (NP_032740.3) (Met1-Gln526) was expressed with the Fc region of human IgG1 at the C-terminus. Predicted N terminal: Ala 19
Species:	Mouse
Expression Host:	CHO Cells
Accession:	Q01705-1
Molecular Weight:	80.6 kDa (predicted)

QC Testing

Biological Activity:	Activity testing is in progress. It is theoretically active, but we cannot guarantee it. If you require protein activity, we recommend choosing the eukaryotic expression version first.
Purity:	> 90 % as determined by SDS-PAGE.
Endotoxin:	< 1.0 EU/ μ g of the protein as determined by the LAL method.
Formulation:	Lyophilized from a solution filtered through a 0.22 μ m filter, containing PBS, pH 7.4. Typically, a mixture containing 5% to 8% trehalose, mannitol, and 0.01% Tween 80 is incorporated as a protective agent before lyophilization.

Preparation and Storage

Reconstitution:

A Certificate of Analysis (CoA) containing reconstitution instructions is included with the products. Please refer to the CoA for detailed information.

Stability & Storage:

It is recommended to store recombinant proteins at -20°C to -80°C for future use. Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Actual storage temperature shall be subject to the COA.

Shipping:

In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

Protein Background

NOTCH1 is one of the four mammalian Notch receptors, which is involved in the Notch signaling pathway. Specifically, NOTCH1 promotes the proliferation of myogenic precursor cells, and the NICD domain of NOTCH1 can impair the regeneration of skeletal muscles. NOTCH1 is a prevalent signaling pathway in T cell acute lymphoblastic leukemia (T-ALL). The NOTCH signaling pathway is a conserved signaling cascade that regulates many aspects of development and homeostasis in multiple organ systems. The proto-oncogene NOTCH1 is frequently mutated in

A DRUG SCREENING EXPERT

around 10% of patients with chronic lymphocytic leukemia (CLL). NOTCH1 mutations in oral squamous cell carcinoma (OSCC) frequently occur near the ligand-binding region. These mutations change the domain structure of this protein and affect the ligand-binding activity. When NOTCH1 is activated by ligand binding, the NOTCH1 intracellular domain (NICD) is cleaved from the cell membrane.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481