

## AGRP Protein, Rhesus, Recombinant (mFc)

### General Information

Synonyms:	agouti related neuropeptide
Protein Construction:	A DNA sequence encoding the rhesus AGRP (XP_001091740.1) (Ser83-Thr132) was expressed with the Fc region of mouse IgG1 at the N-terminus. Predicted N terminal: Asp
Species:	Rhesus
Expression Host:	HEK293 Cells
Accession:	XP_001091740.1
Molecular Weight:	32.3 kDa (predicted)

### QC Testing

Biological Activity:	Activity testing is in progress. It is theoretically active, but we cannot guarantee it. If you require protein activity, we recommend choosing the eukaryotic expression version first.
Purity:	> 90 % as determined by SDS-PAGE
Endotoxin:	< 1.0 EU/µg of the protein as determined by the LAL method.
Formulation:	Lyophilized from a solution filtered through a 0.22 µm filter, containing PBS, pH 7.4. Typically, a mixture containing 5% to 8% trehalose, mannitol, and 0.01% Tween 80 is incorporated as a protective agent before lyophilization.

### Preparation and Storage

**Reconstitution:**  
A Certificate of Analysis (CoA) containing reconstitution instructions is included with the products. Please refer to the CoA for detailed information.

**Stability & Storage:**  
It is recommended to store recombinant proteins at -20°C to -80°C for future use. Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Actual storage temperature shall be subject to the COA.

**Shipping:**  
In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

### Protein Background

Agouti Related Protein (AGRP, or AGRT), is an endogenous antagonist of the melanocortin receptors MC3R and MC4R found in the hypothalamus and exhibits potent orexigenic activity. AGRP can act as a competitive antagonist to proopiomelanocortin (POMC)-derived peptides at the melanocortin-4 receptor (MC4R), and that this homeostatic mechanism is important as a means of coordinating appetite with perceived metabolic requirement. AGRP is upregulated by fasting while intracerebroventricular injections of synthetic AGRP lead to increased

appetite and food intake. Thus, AGRP is a powerful orexigenic peptide that increases food intake when ubiquitously overexpressed or when administered centrally.

### Reference

Ilnytska O, et al. (2008) The role of the Agouti-Related Protein in energy balance regulation. *Cell Mol Life Sci.* 65 (17): 2721-31.

Pritchard LE, et al. (2005) Agouti-related protein: more than a melanocortin-4 receptor antagonist? *Peptides.* 26 (10): 1759-70.

Sttz AM, et al. (2005) The agouti-related protein and its role in energy homeostasis. *Peptides.* 26(10): 1771-81.

Millhauser GL, et al. (2003) Loops and links: structural insights into the remarkable function of the agouti-related protein. *Ann N Y Acad Sci.* 994: 27-35.

**Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins**

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E\_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481