

EMMPRIN/CD147 Protein, Human, Recombinant (hFc)

General Information

| | |
|-----------------------|--|
| Synonyms: | EMMPRIN;CD147;OK;M6;TCSF;Basigin;5F7;basigin (Ok blood group) |
| Protein Construction: | Ala22-His205 |
| Species: | Human |
| Expression Host: | HEK293 Cells |
| Accession: | P35613-2 |
| Molecular Weight: | 47 kDa (predicted); 55-70 kDa (reducing condition, due to glycosylation) |

QC Testing

| | |
|----------------------|--|
| Biological Activity: | Activity testing is in progress. It is theoretically active, but we cannot guarantee it. If you require protein activity, we recommend choosing the eukaryotic expression version first. |
| Purity: | > 95% as determined by SDS-PAGE. |
| Endotoxin: | < 1.0 EU/μg of the protein as determined by the LAL method. |
| Formulation: | Lyophilized from a 0.2 μm filtered solution of 20mM PB, 150mM NaCl, pH 7.2. |

Preparation and Storage

Reconstitution:

Reconstitute the lyophilized protein in distilled water. The product concentration should not be less than 100 μg/ml. Before opening, centrifuge the tube to collect powder at the bottom. After adding the reconstitution buffer, avoid vortexing or pipetting for mixing.

Stability & Storage:

It is recommended to store recombinant proteins at -20°C to -80°C for future use. Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Actual storage temperature shall be subject to the COA.

Shipping:

In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

Protein Background

CD147/EMMPRIN (Extracellular Matrix Metalloproteinase Inducer), also known as Basigin (BSG), is a transmembrane glycoprotein with different forms resulted from different modes of glycosylation and N-terminal sequence variants. It is a member of the immunoglobulin superfamily with homology to both the immunoglobulin V domain and MHC class II antigen beta-chain. This protein play important roles in variety of events including spermatogenesis, embryo implantation, neural network formation. CD147 induces the production and release of matrix metalloproteinases (MMP) in the surrounding mesenchymal cells and tumor cells, and thereby promotes

invasion, metastasis, growth and survival of malignant cells. Furthermore, CD147 also serves as a receptor for extracellular cyclophilin and its association with integrins might be important in signal transduction. Recently, CD147 displays increased expression in many cancers, and it has been previously demonstrated to participate in cancer metastasis and progression. Thus, CD147 and its antibody are used as an effective treatment for malignant cancers.

Reference

- Tang Y, et al. (2004) Tumor-stroma interaction: positive feedback regulation of extracellular matrix metalloproteinase inducer (EMMPRIN) expression and matrix metalloproteinase-dependent generation of soluble EMMPRIN. *Mol Cancer Res.* 2(2): 73-80.
- Wilson MC, et al. (2005) Basigin (CD147) is the target for organomercurial inhibition of monocarboxylate transporter isoforms 1 and 4: the ancillary protein for the insensitive MCT2 is EMBIGIN (gp70). *J Biol Chem.* 280(29): 27213-21.
- Curtin KD, et al. (2005) Basigin (EMMPRIN/CD147) interacts with integrin to affect cellular architecture. *J Cell Sci.* 118 (Pt 12): 2649-60.
- Zhu H, et al. (2009) A novel antibody fragment targeting HAb18G/CD147 with cytotoxicity and decreased immunogenicity. *Cancer Biol Ther.* 8(11): 1035-44.
- Zhu H, et al. (2009) A novel antibody fragment targeting HAb18G/CD147 with cytotoxicity and decreased immunogenicity. *Cancer Biol Ther.* 8(11): 1035-44.
- Seizer P, et al. (2009) EMMPRIN (CD147) is a novel receptor for platelet GPVI and mediates platelet rolling via GPVI-EMMPRIN interaction. *Thromb Haemost.* 101(4): 682-6.
- Moonsom S, et al. (2010) A Competitive ELISA for Quantifying Serum CD147: Reduction of Soluble CD147 Levels in Cancer Patient Sera. *Hybridoma (Larchmt).* 29(1): 45-52.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel: 781-999-4286 E_mail: info@targetmol.com Address: 34 Washington Street, Wellesley Hills, MA 02481