

Nectin-2 Protein, Human, Recombinant (His)

General Information

Synonyms:	PVRR2;Nectin-2;poliovirus receptor-related 2 (herpesvirus entry mediator B);CD112;HVEB; PRR2
Protein Construction:	A DNA sequence encoding the extracellular domain (Met 1-Leu 360) of human CD112 (NP_002847.1) was expressed with a C-terminal polyhistidine tag. Predicted N terminal: Gln 32
Species:	Human
Expression Host:	HEK293 Cells
Accession:	Q92692-2
Molecular Weight:	36.2 kDa (predicted); 48 kDa (reducing condition, due to glycosylation)

QC Testing

Biological Activity:	1. Measured by its binding ability in a functional ELISA. Immobilized recombinant human CD112 at 20 µg/ml (100 µl/well) can bind biotinylated DNAM1 with a linear range of 0.078-2.5 µg/ml. 2. Using the Octet RED System, the affinity constant (Kd) of Anti-CD112 Antibody bound to Nectin-2 Protein, Human, Recombinant (His Tag) was 34.4 nM.
Purity:	≥ 95 % as determined by SDS-PAGE. ≥ 90 % as determined by SEC-HPLC.
Endotoxin:	< 1.0 EU/µg of the protein as determined by the LAL method.
Formulation:	Lyophilized from a solution filtered through a 0.22 µm filter, containing PBS, pH 7.4. Typically, a mixture containing 5% to 8% trehalose, mannitol, and 0.01% Tween 80 is incorporated as a protective agent before lyophilization.

Preparation and Storage

Reconstitution:

Reconstituted with sterile deionized water to 0.25 mg/mL. Reconstitution conditions may vary depending on the lot.

Stability & Storage:

It is recommended to store recombinant proteins at -20°C to -80°C for future use. Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Actual storage temperature shall be subject to the COA.

Shipping:

In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

Protein Background

Cluster of Differentiation 112 (CD112), also known as poliovirus receptor related protein 2 (PVRL2 or PRR2), is a single-pass type I transmembrane glycoprotein belonging to the Immunoglobulin superfamily. CD112 protein also serves as an entry for certain mutant strains of herpes simplex virus and pseudorabies virus, and thus is involved in cell to cell spreading of these viruses. CD112 protein has been identified as the ligand for DNAM-1 (CD226), and the interaction of CD226/CD112 protein can induce NK cell- and CD8+T cell-mediated cytotoxicity and cytokine secretion. CD112 has been regarded as a critical component in allergic reactions, and accordingly may function as a novel target for anti-allergic therapy.

Reference

Bachelet I, et al. (2006) Mast cell costimulation by CD226/CD112 (DNAM-1/Nectin-2): a novel interface in the allergic process. *J Biol Chem.* 281(37): 27190-6.

Wang L, et al. (2009) Molecular cloning, characterization and three-dimensional modeling of porcine nectin-2/CD112. *Vet Immunol Immunopathol.* 132(2-4): 257-63.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481