

EpCAM/TROP1 Protein, Human, Recombinant (hFc)

General Information

Synonyms:	KSA;DIAR5;HNPCC8;TACSTD1;KS1/4;ESA;MK-1;MIC18;epithelial cell adhesion molecule;EGP-2;EGP314;M4S1;TROP-1;TROP1;EGP40
Protein Construction:	A DNA sequence encoding the extracellular domain (Met1-Lys265) of human EpCAM (NP_002345.1) was expressed with the fused Fc region of human IgG1 at the C-terminus. Predicted N terminal: Gln 24
Species:	Human
Expression Host:	HEK293 Cells
Accession:	P16422
Molecular Weight:	54 kDa (predicted); 65-70 kDa (reducing condition, due to glycosylation)

QC Testing

Biological Activity:	Immobilized Recombinant Human EpCAM/TROP-1/TACSTD1 Protein (Fc Tag) (Cat#TMPY-00785) at 1 µg/mL (100 µL/well) can bind anti-EpCAM scFv-hIgG1 Fc, Biotinylated, the EC50 is 0.4-1.5 ng/mL (QC tested).
Purity:	> 95 % as determined by SDS-PAGE
Endotoxin:	< 1.0 EU/µg of the protein as determined by the LAL method.
Formulation:	Lyophilized from a solution filtered through a 0.22 µm filter, containing PBS, pH 7.4. Typically, a mixture containing 5% to 8% trehalose, mannitol, and 0.01% Tween 80 is incorporated as a protective agent before lyophilization.

Preparation and Storage

Reconstitution:

A Certificate of Analysis (CoA) containing reconstitution instructions is included with the products. Please refer to the CoA for detailed information.

Stability & Storage:

It is recommended to store recombinant proteins at -20°C to -80°C for future use. Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Actual storage temperature shall be subject to the COA.

Shipping:

In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

Protein Background

Epithelial Cell Adhesion Molecule (EpCAM), also known as GA733-2 antigen, is a type I transmembrane glycoprotein composed of an extracellular domain with two EGF-Like repeats and a cystenin-rich region, a

transmembrane domain and a cytoplasmic domain. It modulates cell adhesion and proliferation. Its overexpression has been detected in many epithelial tumours and has been associated with high stage, high grade and a worse survival in some tumour types. EpCAM has been shown to function as a calcium-independent homophilic cell adhesion molecule that does not exhibit any obvious relationship to the four known cell adhesion molecule superfamilies. However, recent insights have revealed that EpCAM participates in not only cell adhesion, but also in proliferation, migration and differentiation of cells. In addition, recent study revealed that EpCAM is the Wnt-beta-catenin signaling target gene and may be used to facilitate prognosis. It has oncogenic potential and is activated by release of its intracellular domain, which can signal into the cell nucleus by engagement of elements of the wnt pathway. Cancer Immunotherapy Immune Checkpoint Immunotherapy Targeted Therapy

Reference

Brunner A, et al. (2008) EpCAM is predominantly expressed in high grade and advanced stage urothelial carcinoma of the bladder. *J Clin Pathol.* 61(3):307-10.

Trzpis M, et al. (2008) EpCAM in morphogenesis. *Front Biosci.* 13: 5050-5.

Carpenter G, et al. (2009) EpCAM: another surface-to-nucleus missile. *Cancer Cell.* 15(3): 165-6.

Munz M, et al. (2009) The emerging role of EpCAM in cancer and stem cell signaling. *Cancer Res.* 69(14): 5627-9.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel: 781-999-4286 E_mail: info@targetmol.com Address: 34 Washington Street, Wellesley Hills, MA 02481