

Leptin Protein, Human, Recombinant

General Information

Synonyms:	OBS;leptin;OB;LEPD
Protein Construction:	A DNA sequence encoding the mature form of human Leptin (NP_000221.1) (Val 22-Cys 167) was expressed, with an additional Met. Predicted N terminal: Met
Species:	Human
Expression Host:	E. coli
Accession:	A4D0Y8
Molecular Weight:	16 kDa (predicted); 13 kDa (reducing conditions)

QC Testing

Biological Activity:	Immobilized Human Leptin at 2 µg/mL (100 µL/well) can bind Human Leptin Receptor (His & hFc Tag), the EC50 of Human Leptin Receptor (His & hFc Tag) is 70-420 ng/mL.
Purity:	> 98 % as determined by SDS-PAGE
Endotoxin:	Please contact us for more information.
Formulation:	Lyophilized from a solution filtered through a 0.22 µm filter, containing PBS, pH 7.4. Typically, a mixture containing 5% to 8% trehalose, mannitol, and 0.01% Tween 80 is incorporated as a protective agent before lyophilization.

Preparation and Storage

Reconstitution:
Reconstituted with PBS, pH 7.4 to 1 mg/mL. Reconstitution conditions may vary depending on the lot.

Stability & Storage:

It is recommended to store recombinant proteins at -20°C to -80°C for future use. Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Actual storage temperature shall be subject to the COA.

Shipping:

In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

Protein Background

Leptin is one of the most important hormones secreted by adipocytes, as an adipokine that modulates multiple functions including energy homeostasis, thermoregulation, bone metabolism, endocrine, and pro-inflammatory immune responses. The circulating leptin levels serve as a gauge of energy stores, thereby directing the regulation of energy homeostasis, neuroendocrine function, and metabolism. Recent studies suggest that leptin is physiologically more important as an indicator of energy deficiency, rather than energy excess, and may mediate

adaptation by driving increased food intake and directing neuroendocrine function to conserve energy, such as inducing hypothalamic hypogonadism to prevent fertilization. One of these functions is the connection between nutritional status and immune competence. The adipocyte-derived hormone Leptin has been shown to regulate the immune response, innate, and adaptive response, both in normal and pathological conditions. Thus, Leptin is a mediator of the inflammatory response. Leptin has a dual effect on bone, acting by two independent mechanisms. As a signal molecule with growth factor characteristics, leptin can stimulate osteoblastic cells and inhibit osteoclast formation and activity, thus promoting osteogenesis. However, as a molecule that stimulates sympathetic neurons in the hypothalamus, leptin indirectly inhibits bone formation. This inhibitory effect of leptin mediated by activation of the sympathetic nervous system can be abrogated by the application of blood pressure-reducing beta-blockers, which also inhibit receptors of hypothalamic adrenergic neurons. Leptin appears to regulate some features defining Alzheimer's disease (AD) at the molecular and physiological level. Leptin can stimulate mitogenic and angiogenic processes in peripheral organs. Because leptin levels are elevated in obese individuals and excess body weight has been shown to increase breast cancer risk in postmenopausal women. Furthermore, a recent report clearly shows that targeting leptin signaling may reduce mammary carcinogenesis.

Reference

- Surmacz E. (2007) Obesity hormone leptin: a new target in breast cancer? *Breast Cancer Res.* 9(1): 301.
- Wodarski K, et al. (2009) Leptin as a modulator of osteogenesis. *Ortop Traumatol Rehabil.* 11(1): 1-6.
- Tezapsidis N, et al. (2009) Leptin: a novel therapeutic strategy for Alzheimer's disease. *J Alzheimers Dis.* 16(4): 731-40.
- Cai C, et al. (2009) Leptin in non-autoimmune inflammation. *Inflamm Allergy Drug Targets.* 8(4): 285-91.
- Fernandez-Riejos P, et al. (2010) Role of leptin in the activation of immune cells. *Mediators Inflamm.* 2010: 568343.
- Kelesidis T, et al. (2010) Narrative review: the role of leptin in human physiology: emerging clinical applications. *Ann Intern Med.* 152(2): 93-100.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel: 781-999-4286 E_mail: info@targetmol.com Address: 34 Washington Street, Wellesley Hills, MA 02481