

## Lipocalin-2/LCN2 Protein, Mouse, Recombinant (His)

### General Information

Synonyms:	24p3;lipocalin 2;AW212229;Sip24
Protein Construction:	A DNA sequence encoding the mouse LCN2 (NP_032517.1) precursor (Met 1-Asn 200) was expressed with a C-terminal polyhistidine tag. Predicted N terminal: Gln 20
Species:	Mouse
Expression Host:	HEK293 Cells
Accession:	P11672
Molecular Weight:	22.3 kDa (predicted); 25 kDa (reducing condition, due to glycosylation)

### QC Testing

Biological Activity:	Measured by its ability to bind Iron(III) dihydroxybenzoic acid [Fe(DHBA) <sub>3</sub> ]. The binding of Fe (DHBA) <sub>3</sub> results in the quenching of Trp fluorescence in recombinant mouse Lipocalin-2. Recombinant mouse Lipocalin-2 can bind >1.0 μM of Fe(DHBA) <sub>3</sub> .
Purity:	≥ 90 % as determined by SDS-PAGE
Endotoxin:	< 1.0 EU/μg of the protein as determined by the LAL method.
Formulation:	Lyophilized from a solution filtered through a 0.22 μm filter, containing PBS, pH 7.4. Typically, a mixture containing 5% to 8% trehalose, mannitol, and 0.01% Tween 80 is incorporated as a protective agent before lyophilization.

### Preparation and Storage

**Reconstitution:**  
Reconstituted with sterile deionized water to 0.25 mg/mL. Reconstitution conditions may vary depending on the lot.

**Stability & Storage:**

It is recommended to store recombinant proteins at -20°C to -80°C for future use. Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Actual storage temperature shall be subject to the COA.

**Shipping:**

In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

### Protein Background

Lipocalin-2 (LCN2), also known as neutrophil gelatinase-associated lipocalin (NGAL), is a 25 kDa protein belonging to the lipocalin superfamily. It was initially found in activated neutrophils, however, many other cells, like kidney tubular cells, may produce NGAL in response to various insults. This protein is released from injured tubular cells after various damaging stimuli, is already known by nephrologists as one of the most promising biomarkers of

incoming Acute Kidney Injury (AKI). Recent evidence also suggests its role as a biomarker in a variety of other renal and non-renal conditions. Moreover, recent studies seem to suggest a potential involvement of this factor also in the genesis and progression of chronic kidney diseases. NGAL is the first known mammalian protein that specifically binds organic molecules called siderophores, which are high-affinity iron chelators. NGAL, first known as an antibacterial factor of natural immunity, and an acute-phase protein, is currently one of the most interesting and enigmatic proteins involved in the process of tumor development. acting as an intracellular iron carrier and protecting MMP9 from proteolytic degradation, NGAL has a clear pro-tumoral effect, as has already been observed in different tumors (e.g. breast, stomach, esophagus, brain) in humans. In thyroid carcinomas, NGAL is strongly induced by NF- $\kappa$ B, an important factor involved both in tumor growth and in the link between chronic inflammation and neoplastic development. Thus, Lipocalin-2 (LCN2/NGAL) has been implicated in a variety of processes including cell differentiation, proliferation, survival, and morphogenesis.

### Reference

- Schmidt-Ott KM, et al. (2006) Neutrophil gelatinase-associated lipocalin-mediated iron traffic in kidney epithelia. *Curr Opin Nephrol Hypertens.* 15(4): 442-9.
- Bolignano D, et al. (2010) Neutrophil gelatinase-associated lipocalin (NGAL) in human neoplasias: a new protein enters the scene. *Cancer Lett.* 288(1): 10-6.
- Soni SS, et al. (2010) NGAL: a biomarker of acute kidney injury and other systemic conditions. *Int Urol Nephrol.* 42(1): 141-50.
- Bolignano D, et al. (2010) From kidney to cardiovascular diseases: NGAL as a biomarker beyond the confines of nephrology. *Eur J Clin Invest.* 40(3): 273-6.

**Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins**

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel: 781-999-4286    E\_mail: info@targetmol.com    Address: 34 Washington Street, Wellesley Hills, MA 02481