

FGF-1 Protein, Mouse/Rat, Recombinant

General Information

Synonyms:	fibroblast growth factor 1 (acidic)
Protein Construction:	A DNA sequence encoding the mature form of mouse / rat FGF1 (NP_034327.1) (Phe16-Asp155) was expressed. Mouse and Rat FGF1 sequences are identical. Predicted N terminal: Met
Species:	Mouse,Rat
Expression Host:	E. coli
Accession:	P61148
Molecular Weight:	16 kDa (predicted); 16 kD (reducing conditions)

QC Testing

Biological Activity:	Measured in a cell proliferation assay using Balb/c 3T3 mouse embryonic fibroblast cells. The ED50 for this effect is 0.04-0.2 ng/mL.
Purity:	> 90 % as determined by SDS-PAGE
Endotoxin:	Please contact us for more information.
Formulation:	Lyophilized from a solution filtered through a 0.22 µm filter, containing 50 mM Na2SO4, 0.5 mM DTT, 1 mM EDTA, 20 mM Tris, pH 7.5. Typically, a mixture containing 5% to 8% trehalose, mannitol, and 0.01% Tween 80 is incorporated as a protective agent before lyophilization.

Preparation and Storage

Reconstitution:
Reconstituted with sterile deionized water to 0.25 mg/mL. Reconstitution conditions may vary depending on the lot.

Stability & Storage:

It is recommended to store recombinant proteins at -20°C to -80°C for future use. Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Actual storage temperature shall be subject to the COA.

Shipping:

In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

Protein Background

aFGF, also known as FGF1 and HBGF-1, is a member of the fibroblast growth factor family. The biological activity of aFGF protein is exerted through binding to four high affinity cell surface receptors (FGFR1-4), which results in receptor dimerization and transphosphorylation in the tyrosine kinase domain. aFGF protein shows a wide range of endocrine-like activities. As a multiple function growth factor, this protein is involved in embryo development

and tissue repair. Additionally, this protein is considered to function in several important physiological and pathological processes, such as embryonic development, morphogenesis, angiogenesis, wound healing and atheromatosis, carcinogenesis, development, and invasion of cancer. References

Reference

Jaye M., et al.,(1986), Human endothelial cell growth factor: cloning, nucleotide sequence, and chromosome localization. *Science* 233:541-545.

Mergia A., et al., (1989), Structural analysis of the gene for human acidic fibroblast growth factor. *Biochem. Biophys. Res. Commun.* 164:1121-1129.

Wang W.P., et al.,(1989), Cloning of the gene coding for human class 1 heparin-binding growth factor and its expression in fetal tissues. *Mol. Cell. Biol.* 9:2387-2395.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481