

15-PGDH Protein, Human, Recombinant (His)

General Information

Synonyms:	PGDH1;hydroxyprostaglandin dehydrogenase 15-(NAD);PHOAR1;SDR36C1;15-PGDH;PGDH
Protein Construction:	A DNA sequence encoding the human HPGD (NP_000851.2) (Met 1-Gln 266) was expressed, with a polyhistidine tag at the C-terminus. Predicted N terminal: Met 1
Species:	Human
Expression Host:	E. coli
Accession:	P15428-1
Molecular Weight:	29.7 kDa (predicted); 27 kDa (reducing conditions)

QC Testing

Biological Activity:	Measured by the production of NADH during the oxidation of PGF2 α . The specific activity is >1,500 pmoles/min/ μ g.
Purity:	> 92 % as determined by SDS-PAGE
Endotoxin:	< 1.0 EU/ μ g of the protein as determined by the LAL method.
Formulation:	Supplied as sterile 50 mM Tris, 100 mM NaCl, 0.5 mM DTT, 10% glycerol, pH 7.5.

Preparation and Storage

Reconstitution:

A Certificate of Analysis (CoA) containing reconstitution instructions is included with the products. Please refer to the CoA for detailed information.

Stability & Storage:

It is recommended to store the product under sterile conditions at -20°C to -80°C. Samples are stable for up to 12 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Actual storage temperature shall be subject to the COA.

Shipping:

Proteins are shipped with blue ice.

Protein Background

15-hydroxyprostaglandin dehydrogenase [NAD⁺], also known as Prostaglandin dehydrogenase 1, HPGD, and PGDH1, is a member of the short-chain dehydrogenases/reductases (SDR) family. Prostaglandins (PGs) play a key role in the onset of labor in many species and regulate uterine contractility and cervical dilatation. Therefore, the regulation of prostaglandin output by PG synthesizing and metabolizing enzymes in the human myometrium may determine uterine activity patterns in human labor both at preterm and at term. Prostaglandin dehydrogenase (PGDH) metabolizes prostaglandins (PGs) to render them inactive. HPGD is down-regulated by cortisol, dexamethasone, and betamethasone and down-regulated in colon cancer. It is up-regulated by TGFB1. HPGD contributes to the regulation of events that are under the control of prostaglandin levels. HPGD catalyzes the NAD-

dependent dehydrogenation of lipoxin A4 to form 15-oxo-lipoxin A4. and inhibits in vivo proliferation of colon cancer cells. Defects in HPGD are the cause of primary hypertrophic osteoarthropathy autosomal recessive (PHOAR), cranio-osteoarthropathy (COA), and isolated congenital nail clubbing.

Reference

- Patel, FA. et al., 2003, J. Clin. Endocrinol. Metab. 88: 2922-33.
McKeown KJ, et al.,2003, J. Clin. Endocrinol. Metab. 88 (4): 1737-41.
Yan, M. et al., 2004, Proc. Natl. Acad. Sci. USA. 101: 17468-73.
Tariq, M. et al., 2009, J Med Genet. 46 (1): 14-20.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only· Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481