

HGFR/c-Met Protein, Human, Recombinant (His)

General Information

Synonyms:	AUTS9;MET proto-oncogene, receptor tyrosine kinase;RCCP2;DFNB97;c-Met;HGFR
Protein Construction:	A DNA sequence encoding the extracellular domain of human c-Met (NP_000236.2) (Met1-Thr932) was fused with a polyhistidine tag at the C-terminus. Predicted N terminal: Glu 25
Species:	Human
Expression Host:	HEK293 Cells
Accession:	P08581
Molecular Weight:	103 kDa (predicted); 45 kDa and 85 kDa (reducing condition, due to glycosylation)

QC Testing

Biological Activity:	Immobilized Recombinant Human HGF / Hepatocyte Growth Factor Protein at 2µg/mL (100µL/well) can bind Recombinant Human c-MET / HGFR Protein (His Tag) , the EC50 is 16-48 ng/mL.
Purity:	≥ 90 % as determined by SDS-PAGE. ≥ 90 % as determined by SEC-HPLC.
Endotoxin:	< 1.0 EU/µg of the protein as determined by the LAL method.
Formulation:	Lyophilized from a solution filtered through a 0.22 µm filter, containing 20 mM Tris, 150 mM NaCl, pH8.5. Typically, a mixture containing 5% to 8% trehalose, mannitol, and 0.01% Tween 80 is incorporated as a protective agent before lyophilization.

Preparation and Storage

Reconstitution:

Reconstituted with sterile deionized water to 0.25 mg/mL. Reconstitution conditions may vary depending on the lot.

Stability & Storage:

It is recommended to store recombinant proteins at -20°C to -80°C for future use. Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Actual storage temperature shall be subject to the COA.

Shipping:

In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

Protein Background

Hepatocyte growth factor receptor (HGFR), also known as c-Met or mesenchymal-epithelial transition factor (MET), is a receptor tyrosine kinase (RTK) that is overexpressed and/or mutated in a variety of malignancies. HGFR protein is produced as a single-chain precursor, and HGF is the only known ligand. Normal HGF/HGFR signaling is essential for embryonic development, tissue repair, or wound healing, whereas aberrantly active HGFR has been

strongly implicated in tumorigenesis, particularly in the development of invasive and metastatic phenotypes. HGFR protein is a multifaceted regulator of growth, motility, and invasion, and is normally expressed by cells of epithelial origin. Preclinical studies suggest that targeting aberrant HGFR signaling could be an attractive therapy in cancer. Cancer Immunotherapy Immune Checkpoint Immunotherapy Targeted Therapy

Reference

McGill GG, et al. (2006) c-Met expression is regulated by Mitf in the melanocyte lineage. *J Biol Chem.* 281(15): 10365-73.

Garcia S, et al. (2007) c-Met overexpression in inflammatory breast carcinomas: automated quantification on tissue microarrays. *British journal of cancer.* 96(2): 329-35.

Socoteanu MP, et al. (2008) c-Met targeted therapy of cholangiocarcinoma. *World J Gastroenterol.* 14(19): 2990-4.

Kong DS, et al. (2009) Prognostic significance of c-Met expression in glioblastomas. *Cancer.* 115(1): 140-8.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel: 781-999-4286 E_mail: info@targetmol.com Address: 34 Washington Street, Wellesley Hills, MA 02481