

## Carboxypeptidase B1 Protein, Mouse, Recombinant (His)

## General Information

Synonyms:	carboxypeptidase B1 (tissue);1810063F02Rik;AI504870;0910001A18Rik;2210008M23Rik
Protein Construction:	A DNA sequence encoding the mouse CPB1 (NP_083982.1) precursor (Met 1-Tyr 415) was expressed, with a C-terminal polyhistidine tag. Predicted N terminal: Ser 16
Species:	Mouse
Expression Host:	HEK293 Cells
Accession:	B2RS76
Molecular Weight:	47.5 kDa (predicted); 43-45 kDa (reducing conditions)

## QC Testing

Biological Activity:	Measured by its ability to cleave a colorimetric peptide substrate, Hippuryl-Arg. The specific activity is >5,000 pmoles/min/ $\mu$ g.
Purity:	> 94 % as determined by SDS-PAGE
Endotoxin:	< 1.0 EU/ $\mu$ g of the protein as determined by the LAL method.
Formulation:	Lyophilized from a solution filtered through a 0.22 $\mu$ m filter, containing 20 mM MES, 150 mM NaCl, pH 6.5. Typically, a mixture containing 5% to 8% trehalose, mannitol, and 0.01% Tween 80 is incorporated as a protective agent before lyophilization.

## Preparation and Storage

## Reconstitution:

A Certificate of Analysis (CoA) containing reconstitution instructions is included with the products. Please refer to the CoA for detailed information.

## Stability &amp; Storage:

It is recommended to store recombinant proteins at -20°C to -80°C for future use. Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Actual storage temperature shall be subject to the COA.

## Shipping:

In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

## Protein Background

Carboxypeptidase B1, also well known as pancreatic procarboxypeptidase B (PCPB), is a highly pancreas -specific protein (PASP), and has been identified previously as a serum marker for acute pancreatitis and pancreatic graft rejection. As the prototype for those human exopeptidases that cleave off basic C-terminal residues, CPB1 specifically cleaves the C-terminal Arg and Lys residues with a preference for Arg. The B1 and B2 forms of procarboxypeptidase B differ from each other mainly in isoelectric point. The deduced amino acid sequence of

## A DRUG SCREENING EXPERT

---

PCPB predicts a 416-amino acid preproenzyme consisting of a 15-aa signal peptide, a 95-aa activation peptide and a 37-aa mature chain. The secreted PCPB zymogen is converted to enzymatically active CPB1 by limited proteolysis by trypsin.

### Reference

Yamamoto, K.K. et al., 1992, J. Biol. Chem. 267: 2575-2581.

Pezzilli, R. et al., 1994, Digestion. 55: 73-77.

Barbosa Pereira, P.J. et al., 2002, J. Mol. Biol. 321: 537-547.

**Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins**

This product is for Research Use Only· Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E\_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481