

IL-12RB1 Protein, Human, Recombinant (His)

General Information

Synonyms:	interleukin 12 receptor, β 1; IL12RB; IMD30; IL-12R-BETA1; interleukin 12 receptor, beta 1; CD212; IL-12R- β 1
Protein Construction:	A DNA sequence encoding the human IL12RB1 isoform 1 (NP_005526.1) extracellular domain (Met1-Glu540) was expressed fused with a polyhistidine tag at the C-terminus. Predicted N terminal: Cys 24
Species:	Human
Expression Host:	HEK293 Cells
Accession:	P42701-1
Molecular Weight:	58.5 kDa (predicted); 70 kDa (reducing condition, due to glycosylation)

QC Testing

Biological Activity:	Measured by its binding ability in a functional ELISA. Immobilized human IL12RB1 at 20 μ g/ml (100 μ l/well) can bind human IL12B with a linear range of 2.56-64 ng/ml.
Purity:	> 94 % as determined by SDS-PAGE
Endotoxin:	< 1.0 EU/ μ g of the protein as determined by the LAL method.
Formulation:	Lyophilized from a solution filtered through a 0.22 μ m filter, containing PBS, pH 7.4. Typically, a mixture containing 5% to 8% trehalose, mannitol, and 0.01% Tween 80 is incorporated as a protective agent before lyophilization.

Preparation and Storage

Reconstitution:	Reconstituted with sterile deionized water to 0.25 mg/mL. Reconstitution conditions may vary depending on the lot.
Stability & Storage:	It is recommended to store recombinant proteins at -20°C to -80°C for future use. Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots. <small>Actual storage temperature shall be subject to the COA.</small>
Shipping:	In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

Protein Background

Interleukin 12 receptor, beta 1 is also known as IL-12 receptor beta component, IL-12R subunit beta-1, and CD212 antigen (CD212). IL12RB1(CD212) is a subunit of the interleukin 12 receptor. IL12RB1(CD212) is a type I transmembrane protein that belongs to the hemopoietin receptor superfamily. This protein binds to interleukin 12

(IL12) with a low affinity and is thought to be a part of the IL12 receptor complex. IL12RB1(CD212) forms a disulfide-linked oligomer, which is required for its IL12 binding activity. The coexpression of IL12RB1 and IL12RB2 proteins was shown to lead to the formation of high-affinity IL12 binding sites and reconstitution of IL12 dependent signaling. The lack of expression of this gene was found to result in the immunodeficiency of patients with severe mycobacterial and Salmonella infections. IL12RB1(CD212) Functions as an interleukin receptor that binds interleukin-12 with low affinity and is involved in IL12 transduction. It is associated with IL12RB2 it forms a functional, high-affinity receptor for IL12. IL12RB1(CD212) associates also with IL23R to form the interleukin-23 receptor which functions in IL23 signal transduction probably through activation of the Jak-Stat signaling cascade.

Reference

Cleary AM, et al. (2003) Impaired accumulation and function of memory CD4 T cells in human IL-12 receptor beta 1 deficiency. *J Immunol.* 170 (1): 597-603.

Suzuki Y, et al. (1997) Construction and characterization of a full length-enriched and a 5'-end-enriched cDNA library. *Gene.* 200 (1-2): 149-56.

Yamamoto K, et al. (1997) Assignment of IL12RB1 and IL12RB2, interleukin-12 receptor beta 1 and beta 2 chains, to human chromosome 19 band p13.1 and chromosome 1 band p31.2, respectively, by in situ hybridization.

Cytogenet. Cell Genet. 77 (3-4): 257-8.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481