

CXADR/CAR Protein, Mouse, Recombinant (His)

General Information

Synonyms:	CAR;coxsackie virus and adenovirus receptor;MCAR;AU016810;MCVADR;2610206D03Rik;AW553441
Protein Construction:	A DNA sequence encoding the mouse CXADR (NP_001020363.1) extracellular domain (Met 1-Gly 237) was fused with a polyhistidine tag at the C-terminus. Predicted N terminal: Leu 20
Species:	Mouse
Expression Host:	HEK293 Cells
Accession:	P97792-1
Molecular Weight:	25.7 kDa (predicted); 35 kDa (reducing condition, due to glycosylation)

QC Testing

Biological Activity:	Activity testing is in progress. It is theoretically active, but we cannot guarantee it. If you require protein activity, we recommend choosing the eukaryotic expression version first.
Purity:	> 95 % as determined by SDS-PAGE
Endotoxin:	< 1.0 EU/μg of the protein as determined by the LAL method.
Formulation:	Lyophilized from a solution filtered through a 0.22 μm filter, containing PBS, pH 7.4. Typically, a mixture containing 5% to 8% trehalose, mannitol, and 0.01% Tween 80 is incorporated as a protective agent before lyophilization.

Preparation and Storage

Reconstitution:
Reconstituted with sterile deionized water to 0.25 mg/mL. Reconstitution conditions may vary depending on the lot.

Stability & Storage:

It is recommended to store recombinant proteins at -20°C to -80°C for future use. Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Actual storage temperature shall be subject to the COA.

Shipping:

In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

Protein Background

CXADR (coxsackie virus and adenovirus receptor), also known as CAR, is a type I transmembrane glycoprotein belonging to the CTX family of the Ig superfamily, and is essential for normal cardiac development in the mouse. Proposed as a homophilic cell adhesion molecule, CXADR is a component of the epithelial apical junction complex that is essential for the tight junction integrity, and probably involved in transepithelial migration of

polymorphonuclear leukocytes (PMN). Mature mouse CXADR structurally comprises a 218 aa extracellular domain (ECD) with a V-type (D1) and a C2-type (D2) Ig-like domain, a 21 aa transmembrane segment and a 17 aa intracellular domain, among which, D1 is thought to be responsible for homodimer formation in trans within tight junctions. The ECD of mouse CXADR shares 97%, 9% sequence identity with the corresponding regions of rat, human CXADR.

Reference

Tomko, R.P. et al., 1997, Proc. Natl. Acad. Sci. U.S.A. 94 (7): 3352-3356.

van Raaij, M.J. et al., 2001, Structure. 8 (11): 1147-1155.

Cohen, C.J. et al., 2001, J. Biol. Chem. 276 (27): 25392-25398.

Carson, S.D. et al., 2002, Rev. Med. Virol. 11 (4): 219-226.

Selinka, H.C. et al., 2004, Med. Microbiol. Immunol. 193 (2-3): 127-131.

Philipson, L. et al., 2004, Curr. Top. Microbiol. Immunol. 273:87-111.

Raschperger, E. et al., 2006, Exp. Cell Res. 312: 1566-1580.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481