

## HER2/ERBB2 Protein, Rhesus, Recombinant (His)

### General Information

Synonyms:	EGFR2;erb-b2 receptor tyrosine kinase 2
Protein Construction:	A DNA sequence encoding the Rhesus ErbB2 (XP_001090430.1) extracellular domain (Met 1-Thr 652) was fused with a polyhistidine tag at the C-terminus. Predicted N terminal: Gln 24
Species:	Rhesus
Expression Host:	HEK293 Cells
Accession:	XP_001090430.1
Molecular Weight:	70.7 kDa (predicted); 105 kDa (reducing condition, due to glycosylation)

### QC Testing

Biological Activity:	Measured by its binding ability in a functional ELISA. Immobilized Rhesus Her2/ERBB2 His at 2 µg/ml (100 µl/well) can bind Herceptin, the EC50 of Herceptin is 10-50 ng/mL.
Purity:	> 95 % as determined by SDS-PAGE. > 90 % as determined by SEC-HPLC.
Endotoxin:	< 1.0 EU/µg of the protein as determined by the LAL method.
Formulation:	Lyophilized from a solution filtered through a 0.22 µm filter, containing PBS, pH 7.4. Typically, a mixture containing 5% to 8% trehalose, mannitol, and 0.01% Tween 80 is incorporated as a protective agent before lyophilization.

### Preparation and Storage

**Reconstitution:**  
A Certificate of Analysis (CoA) containing reconstitution instructions is included with the products. Please refer to the CoA for detailed information.

**Stability & Storage:**

It is recommended to store recombinant proteins at -20°C to -80°C for future use. Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Actual storage temperature shall be subject to the COA.

**Shipping:**

In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

### Protein Background

Human epidermal growth factor receptor 2 (HER2), also known as ErbB2, NEU, and CD340, is a type I membrane glycoprotein and belongs to the epidermal growth factor (EGF) receptor family. HER2 protein cannot bind growth factors due to the lacking of ligand binding domain of its own and autoinhibited constitutively. However, HER2 forms a heterodimer with other ligand-bound EGF receptor family members, therefore stabilizes ligand binding and enhances kinase-mediated activation of downstream molecules. HER2 plays a key role in development, cell

proliferation and differentiation. HER2 gene has been reported to associate with malignancy and a poor prognosis in numerous carcinomas, including breast, prostate, ovarian, lung cancers and so on. Cancer Immunotherapy Immune Checkpoint Immunotherapy Targeted Therapy

### Reference

Krawczyk N, et al. (2009) HER2 status on persistent disseminated tumor cells after adjuvant therapy may differ from initial HER2 status on primary tumor. *Anticancer Res.* 29(10): 4019-24.

**Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins**

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel: 781-999-4286 E\_mail: info@targetmol.com Address: 34 Washington Street, Wellesley Hills, MA 02481