

GRO beta/CXCL2 Protein, Mouse, Recombinant (His & SUMO)

General Information

Synonyms:	Mip2; GRO β /CXCL2; chemokine (C-X-C motif) ligand 2; GRO β ; MIP-2; Scyb2; Mgsa-b; Scyb; Gro2; MIP-2a; CINC-2a
Protein Construction:	A DNA sequence encoding the mature form of mouse CXCL2 (P10889-1) (Ala 28-Asn 100) was fused with a N-terminal polyhistidine-tagged SUMO tag at the N-terminus. Predicted N terminal: Met
Species:	Mouse
Expression Host:	E. coli
Accession:	P10889-1
Molecular Weight:	21.2 kDa (predicted); 30 kDa (reducing conditions)

QC Testing

Biological Activity:	Activity testing is in progress. It is theoretically active, but we cannot guarantee it. If you require protein activity, we recommend choosing the eukaryotic expression version first.
Purity:	> 95 % as determined by SDS-PAGE
Endotoxin:	Please contact us for more information.
Formulation:	Lyophilized from a solution filtered through a 0.22 μ m filter, containing 0.1% TFA, pH 2.9. Typically, a mixture containing 5% to 8% trehalose, mannitol, and 0.01% Tween 80 is incorporated as a protective agent before lyophilization.

Preparation and Storage

Reconstitution:

A Certificate of Analysis (CoA) containing reconstitution instructions is included with the products. Please refer to the CoA for detailed information.

Stability & Storage:

It is recommended to store recombinant proteins at -20°C to -80°C for future use. Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Actual storage temperature shall be subject to the COA.

Shipping:

In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

Protein Background

Chemokine (C-X-C motif) ligand 2 (CXCL2), also called macrophage inflammatory protein 2 (MIP-2), Growth-regulated protein beta (Gro-beta) and Gro oncogene-2 (Gro-2), is a small cytokine belonging to the CXC chemokine family. CXCL2/MIP-2 is selectively up-regulated in tolerance-conferring APCs and serves to recruit NKT

cells to the splenic marginal zone, where they form clusters with APCs and T cells. In the absence of the high-affinity receptor for CXCL2/MIP-2 or in the presence of a blocking Ab to CXCL2/MIP-2, peripheral tolerance is prevented, and Ag-specific T regulatory cells are not generated. CXCL2/MIP-2 is selectively up-regulated in tolerance-conferring APCs and serves to recruit NKT cells to the splenic marginal zone, where they form clusters with APCs and T cells. In the absence of the high-affinity receptor for MIP-2 (as in CXCR2-deficient mice) or in the presence of a blocking Ab to MIP-2, peripheral tolerance is prevented, and Ag-specific T regulatory cells are not generated. Understanding the regulation of lymphocyte traffic during tolerance induction may lead to novel therapies for autoimmunity, graft acceptance, and tumor rejection. Several studies have implicated the CXCL2 chemokine as a mediator in the development of sepsis. CXCL2/MIP-2 also plays a major role in mediating the neutrophilic inflammatory response of the rodent lung to particles such as quartz, crocidolite asbestos, as well as high doses of other relative innocuous dusts such as titanium dioxide.

Reference

- Driscoll KE. (2000) TNFalpha and MIP-2: role in particle-induced inflammation and regulation by oxidative stress. *Toxicol Lett.* 112-113: 177-83.
- Walpen S, et al. (2001) Nitric oxide induces MIP-2 transcription in rat renal mesangial cells and in a rat model of glomerulonephritis. *FASEB J.* 15(3): 571-3.
- Fahey TJ, et al. (1990) Cytokine production in a model of wound healing: the appearance of MIP-1, MIP-2, cachectin/TNF and IL-Cytokine. *J Biol Chem.* 265(1): 92-9.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481